

**LONG ACTING COUGH SOFTGELS- dextromethorphan hydrobromide capsule,  
liquid filled**

**Chain Drug Marketing Association, Inc.**

-----  
**QC® QUALITY CHOICE Adult Long Acting Cough Softgels**

***Drug Facts***

***Active ingredient (in each softgel)***

Dextromethorphan HBr, USP 15 mg

***Purpose***

Cough Suppressant

***Use***

temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold.

***Warnings***

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a cough that occurs with too much phlegm (mucus)
- a cough that lasts or is chronic as occurs with smoking, asthma, or emphysema

**Stop use and ask a doctor if** cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These can be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

***Directions***

do not take more than 8 softgels in any 24-hour period

this adult product is not intended for use in children under 12 years of age

adults and children 12 years and older	take 2 softgels every 6 to 8 hours, as needed
children under 12 years	do not use

### ***Other information***

- store at 20°-25°C(68°-77°F)
- avoid excessive heat above 40°C(104°F)
- protect from light

### ***Inactive ingredients***

edible white ink, FD&C blue #1, FD&C red #40, gelatin, glycerin, isopropyl alcohol, medium chain triglycerides, polyethylene glycol, povidone, propyl gallate, propylene glycol, purified water, sorbitol sorbitan solution

### ***Questions or comments?***

**1-888-577-8033** Monday - Friday 8am - 4pm EST

### **Compare to the active ingredient in Robitussin® CoughGels®\***

Relieves cough for up to 8 Hours  
Non-Drowsy

### **READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

### **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING FROM BOTTLE**

Distributed by: C.D.M.A., Inc.©  
43157 W 9 Mile Rd Novi, MI 48375  
www.qualitychoice.com  
Product of UAE

Packaged and Quality Assured in the USA

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Robitussin® CoughGels®.

### **100% QC SATISFACTION GUARANTEED**

### **Packaging**

**INNER PACKAGE LABEL**

**QC QUALITY CHOICE** NDC 63868-824-20

Compare to the active ingredient in Robitussin® CoughGels\*\*

# Adult Long Acting Cough Softgels

Dextromethorphan HBr, USP 15 mg

Cough Suppressant  
Relieves cough for up to 8 Hours  
Non-Drowsy

**20 SOFTGELS** Actual Size

**READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**Drug Facts**

**Purpose**  
Dextromethorphan HBr, USP 15 mg,.....Cough Suppressant

**Use**  
Temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold.

**Warnings**  
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** a cough that occurs with too much phlegm (mucus) a cough that lasts or is chronic as occurs with smoking, asthma, or emphysema

**Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These can be signs of a serious condition.**

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions** do not take more than 8 softgels in any 24-hour period this adult product is not intended for use in children under 12 years of age adults and children 12 years and older: take 2 softgels every 6 to 8 hours, as needed as needed children under 12 years: do not use

**TAMPER EVIDENT DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING FROM BOTTLE**

Product of UAE  
Packaged and Quality Assured in the USA  
REV.00-072021  
LB6396882420

LOT:  
EXP:

**No Varnish**

**OUTER PACKAGE LABEL**

**QC QUALITY CHOICE** NDC 63868-824-20

Compare to the active ingredient in Robitussin® CoughGels\*\*

# Adult Long Acting Cough Softgels

Dextromethorphan HBr, USP 15 mg

Cough Suppressant  
Relieves cough for up to 8 Hours  
Non-Drowsy

**20 SOFTGELS** Actual Size

**READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**Warnings (continued)**  
critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**  
do not take more than 8 softgels in any 24-hour period  
this adult product is not intended for use in children under 12 years of age

adults and children 12 years and older	take 2 softgels every 6 to 8 hours, as needed
children under 12 years	do not use

**Other information**  
store at 20°-25°C (68°-77°F)  
avoid excessive heat above 40° C (104° F)  
protect from light

**Inactive ingredients**  
edible white ink, FD&C blue #1, FD&C red #40, gelatin, glycerin, isopropyl alcohol, medium chain triglycerides, polyethylene glycol, povidone, propyl gallate, propylene glycol, purified water, sorbitol sorbitan solution

**Questions or comments?**  
1-888-577-8033 Monday-Friday  
8am-4pm EST

**Ingredients**  
Dextromethorphan HBr, USP 15 mg,.....Cough Suppressant

relieves cough due to minor thro at and irritation as may occur with a cold.

**Warnings**  
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or disease), or for 2 weeks after stopping the MAOI, ask a doctor or pharmacist before use.

**Ask a doctor before use if you have** a cough that occurs with too much phlegm (mucus) that lasts or is chronic as occurs with asthma, or emphysema

**Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These can be signs of a serious condition.**

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions** do not take more than 8 softgels in any 24-hour period this adult product is not intended for use in children under 12 years of age adults and children 12 years and older: take 2 softgels every 6 to 8 hours, as needed as needed children under 12 years: do not use

**TAMPER EVIDENT DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING FROM BOTTLE**

Product of UAE  
Packaged and Quality Assured in the USA  
REV.00-072021  
CT6396882420

LOT:  
EXP:

**No Varnish**

6 35515 99612 0

100% SATISFACTION GUARANTEE  
Distributed by: Q.D.M.A., Inc. @ 43157 W. 9 Mile, Bel Air, MD 43875 www.qualitychoice.com Product of UAE  
Packaged and Quality Assured in the USA  
This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Robitussin® CoughGels®.



**DRUG FACTS TABLE**

<p><b>Drug Facts</b></p> <table border="1"> <tr> <td style="width: 60%;"><b>Active ingredient (in each softgel)</b></td> <td style="width: 40%;"><b>Purpose</b></td> </tr> <tr> <td>Dextromethorphan HBr, USP 15 mg.....</td> <td>Cough Suppressant</td> </tr> </table> <p><b>Use</b> temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold.</p> <p><b>Warnings</b> Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>■ a cough that occurs with too much phlegm (mucus)</li> <li>■ a cough that lasts or is chronic as occurs with smoking, asthma, or emphysema</li> </ul> <p><b>Stop use and ask a doctor if</b> cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These can be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use.</p> <p><b>Keep out of reach of children. Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is</p>	<b>Active ingredient (in each softgel)</b>	<b>Purpose</b>	Dextromethorphan HBr, USP 15 mg.....	Cough Suppressant	<p><b>Warnings (continued)</b> critical for adults as well as children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ do not take more than 8 softgels in any 24-hour period</li> <li>■ this adult product is not intended for use in children under 12 years of age</li> </ul> <table border="1"> <tr> <td>adults and children 12 years and older</td> <td>take 2 softgels every 6 to 8 hours, as needed</td> </tr> <tr> <td>children under 12 years</td> <td>do not use</td> </tr> </table> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store at 20°-25°C(68°-77°F)</li> <li>■ avoid excessive heat above 40°C(104°F)</li> <li>■ protect from light</li> </ul> <p><b>Inactive ingredients</b> edible white ink, FD&amp;C blue #1, FD&amp;C red #40, gelatin, glycerin, isopropyl alcohol, medium chain triglycerides, polyethylene glycol, povidone, propyl gallate, propylene glycol, purified water, sorbitol sorbitan solution</p> <p><b>Questions or comments?</b> 1-888-577-8033 Monday-Friday 8am-4pm EST</p>	adults and children 12 years and older	take 2 softgels every 6 to 8 hours, as needed	children under 12 years	do not use
<b>Active ingredient (in each softgel)</b>	<b>Purpose</b>								
Dextromethorphan HBr, USP 15 mg.....	Cough Suppressant								
adults and children 12 years and older	take 2 softgels every 6 to 8 hours, as needed								
children under 12 years	do not use								

**LONG ACTING COUGH SOFTGELS**

dextromethorphan hydrobromide capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-824
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name		Basis of Strength	Strength	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	15 mg	
<b>Inactive Ingredients</b>				
Ingredient Name		Strength		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)				
GLYCERIN (UNII: PDC6A3C0OX)				
ISOPROPYL ALCOHOL (UNII: ND2M416302)				
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
PROPYL GALLATE (UNII: 8D4SNN7V92)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SORBITOL (UNII: 506T60A25R)				
SORBITAN (UNII: 6O921CV9RU)				
<b>Product Characteristics</b>				
Color	red	Score	no score	
Shape	OVAL (oblong)	Size	14mm	
Flavor		Imprint Code	778	
Contains				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-824-20	1 in 1 CARTON	05/03/2022	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	05/03/2022		

**Labeler** - Chain Drug Marketing Association, Inc. (011920774)