

**TUKOL MAX ACTION SEVERE CONGESTION AND COUGH- dextromethorphan
hbr, guaifenesin, phenylephrine hcl liquid
Genomma Lab USA, Inc**

Tukol MAX ACTION Severe Congestion & Cough

Drug Facts

**Active ingredients
(in each 20 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCL 10 mg

Purpose

Cough suppressant

Expectorant

Nasal decongestant

Uses

- Helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping an MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

taking any other oral nasal decongestant or stimulant.

Do not exceed recommended dosage.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a professional before use

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter

age	dose
adults and children 12 years of age and older	20 mL every 4 hours
children under 12 years of age	do not use

Other information

- Each 20 mL dose contains: sodium 10 mg
- store between 15-30°C (59-86 °F)
- do not refrigerate

Inactive Ingredients

Anhydrous citric acid, edetate disodium, FD&C Blue # 1, FD&C Red # 40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-877-994-3666

Monday to Friday from 8 AM to 6 PM, Central time

Tukol MAX ACTION Severe Congestion & Cough

Tukol[®]

MAX ACTION

DO NOT USE IF PRINTED SEAL UNDER CAP IS
TORN OR MISSING / NO SE USE SI EL SELLO IMPRESO
DEBAJO DE LA TAPA ESTA ROTO O FALTA

Tukol[®]

MAX ACTION

SEVERE CONGESTION & COUGH
CONGESTIÓN Y TOS SEVERA

Dextromethorphan HBr / Guaifenesin / Phenylephrine HCl
Dextrometorfano HBr / Guaifenesina / Fenilefrina HCl

Ages / Edades
12+

MAX
STRENGTH/
MÁXIMA
POTENCIA

Relieves / Alivia:

- **COUGH / TOS**
- **NASAL CONGESTION / CONGESTIÓN NASAL**
- **CHEST CONGESTION / CONGESTIÓN DE PECHO**

6 FL OZ (177 mL)



Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter

age	dose
adults and children 12 years of age and older	20 mL every 4 hours
children under 12 years of age	do not use

Other information

- each (20 mL) dose contains: sodium 10 mg
- store between 15-30°C (59-86°F) do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

Questions or comments? 1-877-994-3666
Monday to Friday from 8 am to 6 pm, Central Time.

Información del Medicamento

Ingredientes activos (en cada 20 mL)

Dextrometorfano HBr, 20 mg.....Inhibidor de la tos
Guaifenesina, 400 mg.....Expectorante
Fenilefrina HCl, 10 mg.....Descongestionante nasal

Propósitos

Usos

- ayuda a desprender las flemas (mucosidad) y a adelgazar las secreciones bronquiales para drenar las vías respiratorias
- alivia temporalmente los siguientes síntomas que se presentan con el resfriado:
 - congestión nasal
 - tos debida a irritación leve de garganta y bronquios.

Advertencias

No utilizarlo si usted está tomando un inhibidor de la monoamina oxidasa (IMAO) de prescripción (ciertos medicamentos para la depresión, condiciones psiquiátricas o emocionales, o enfermedad de Parkinson), o durante 2 semanas después de suspender el medicamento IMAO. Si usted desconoce si su medicamento de prescripción contiene un IMAO, consulte a su médico o farmacéutico antes de tomar este producto.

Consulte a su médico antes de usar, si padece

- enfermedad cardíaca
- hipertensión
- enfermedad de la tiroides
- diabetes
- dificultad al orinar debido a dilatación de la glándula prostática
- tos que se presenta con exceso de flema (mucosidad)
- tos persistente o crónica tal como ocurre con el tabaquismo, asma, bronquitis crónica o enfisema.

Información del Medicamento (continuación)

Consulte a su médico o farmacéutico antes de utilizarlo si usted está tomando cualquier otro descongestionante o estimulante nasal oral.

No exceda la dosis recomendada.

- Suspenda su uso y consulte a su médico si
 - se siente nervioso, mareado o con insomnio
 - los síntomas no mejoran en los siguientes 7 días o se acompañan de fiebre
 - la tos persiste por más de 7 días, regresa, o se presenta acompañada por fiebre, erupción o dolor de cabeza persistente. Estos podrían ser signos de una condición grave.

En caso de embarazo o lactancia, consulte a un profesional médico antes de su uso.

Manténgase fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o contacte al Centro de Control de Envenenamiento de manera inmediata.

Indicaciones

- no tomar más de 6 dosis en un período de 24 horas
- este producto de concentración para adultos no está previsto para su uso en niños menores de 12 años de edad
- medir solo con la copa dosificadora incluida
- mantenga la copa dosificadora con este producto
- mL = mililitro

edad	dosis
adultos y niños de 12 años de edad y mayores	20 mL cada 4 horas
niños menores de 12 años	no usar

Otra información

- cada (20 mL) contiene: sodio 10 mg
- almacenar entre 15 y 30°C (59 a 86°F)
- no refrigerar.

Ingredientes inactivos

ácido cítrico anhidro, edetato disódico, FD&C azul #1, FD&C rojo #40, sabores, glicerina, propilenglicol, galato de propilo, agua purificada, benzoato sódico, sorbitol, sucralosa, goma xantana.

¿Preguntas o comentarios?

1-877-994-3666 Lunes a Viernes de 8 am a 6 pm, Hora del Centro.

Drug Facts

Active ingredients Purposes (in each 20 mL)

Dextromethorphan HBr, 20 mg.....Cough suppressant
Guaifenesin, 400 mg.....Expectorant
Phenylephrine HCl, 10 mg.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

Do not exceed recommended dosage.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Distributed by / Distribuido por:
Genomma Lab USA Inc.,
Houston, TX 77098
BX-064

2000004535 V-3



LOT No. / LOTE
EXP.

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

DEXTROMETHORPHAN HBr/GUAIFENESIN/PHENYLEPHRINE HCL

Relieves

- Cough
- Nasal Congestion
- Chest Congestion

Ages 12+

6 FL OZ (177 mL)

Distributed by:
Genomma Lab USA, Inc.
Houston, TX 77098

TUKOL MAX ACTION SEVERE CONGESTION AND COUGH

dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50066-517
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL GALLATE (UNII: 8D45NN7V92)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-517-25	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/16/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/16/2016	

Labeler - Genomma Lab USA, Inc (832323534)

Revised: 10/2024

Genomma Lab USA, Inc