

**MAX COLD, FLU, AND SORE THROAT- acetaminophen,dextromethorphan hydrobromide, guaifenesin , phenylephrine hcl liquid  
H E B**

-----  
**Drug Facts Active ingredients**

**Active ingredients (in each 20 mL)**

Acetaminophen 650 mg  
Dextromethorphan HBr 20 mg  
Guaifenesin 400 mg  
Phenylephrine HCL 10 mg

**Purpose**

Pain reliever/Fever reducer  
Cough suppressant  
Expectorant  
Nasal Decongestant

**Keep out of reach of children.**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Uses**

temporarily relieves these common cold and flu symptoms:

- cough due to minor throat and bronchial irritation
- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- sore throat
- headache
- fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**

**Liver warning** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reaction. Symptoms may include:

- skin reddening - blisters ■ rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- for children under 12 years of age

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin.

### **When using this product**

- **do not use more than directed**

### **Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever

- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

### **If pregnant or breast feeding**

ask a health professional before use.

### **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact the Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

Age	Dose
Adults and children 12 years and older	20 mL every 4 hours
Children under 12 years of age	Do not use

### **Other information**

- each 20 mL contains: sodium 10 mg
- dosage cup provided
- store between 20°-25° C (68°-77° F)
- do not refrigerate

### **Inactive ingredients**

Anhydrous citric acid, edetate disodium, FD and C Blue #1, FD and C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, Xanthan gum

### **Questions?**

Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

### **Product Label**

**Compare to Mucinex® FAST-MAX™**

**Cold, Flu & Sore Throat** active ingredients\*

NDC 37808-316-25

**H E B®**

**Maximum Strength**

**MAX COLD, FLU & SORE THROAT**

**Acetaminophen /**

(Pain Reliever/Fever Reducer)

Dextromethorphan HBr /

(Cough Suppressant)

Guaifensin / (Expectorant)

Phenylephrine HCL /

Nasal Decongestant)

**Cold, Flu & Throat**

Relief of:

- Fever and Body aches
- Controls Cough
- Thins and Loosens Mucus
- Relieves Nasal and Chest Congestion

**Adults**

For ages 12 and Over

**6 FL OZ (117 mL)**

LF-002 Rev.2 02

**Peel Corner to Read Complete Drug Facts and Information**

**DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING**

**Failure to follow these warnings could result in consequences**

100%

GUARANTEE

promise

If you aren't completely pleased  
with this product, we'll be happy to  
replace it or refund your money.

You have our word on it.

LB-002 Rev 02

1208

Lot:

Exp:

Compare to Mucinex® FAST-MAX™  
Cold, Flu & Sore Throat active ingredients\*

NDC 37808-316-25



Maximum Strength  
**Max Cold, Flu  
& Sore Throat**

Acetaminophen /  
Pain Reliever/Fever Reducer  
Dextromethorphan HBr /  
Cough Suppressant  
Guaifenesin / Expectorant  
Phenylephrine HCl /  
Nasal Decongestant

**Cold, Flu & Throat**

Relief of:

- Fever & Body Aches
- Cough
- Thins & Loosens Mucus
- Nasal & Chest Congestion

**Adults**

For Ages 12 & Over

**6 FL OZ (177mL)**

LF-002 Rev. 02

Peel Corner to Read Complete Drug Facts and Information



DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Failure to follow these warnings could result in serious consequences



100% GUARANTEE promise

If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

LB-002 Rev 02 1208

8581-1711



0 41220 38701 4

Lot:

Exp:

### Drug Facts

#### Active ingredients Purpose (in each 20 mL)

Acetaminophen 650 mg...	Pain reliever/ Fever reducer
Dextromethorphan HBr 20 mg....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

**Uses** temporarily relieves these common cold and flu symptoms:

- cough due to minor throat and bronchial irritation
- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- sore throat
- headache
- fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

#### Warnings

**Liver warning** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reaction. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away. ▶

### Drug Facts (continued)

**Sore throat warning** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

**Do not use ■ with any other drug containing acetaminophen** (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ for children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus) ▶

**Drug Facts** (continued)

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin.

When using this product ■ do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Overdose warning** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even - if you do not notice any signs or symptoms. ►

**Drug Facts** (continued)**Directions**

- take only as recommended (see Overdose warning)
- use dosage cup ■ mL = milliliter
- do not take more than 6 doses in any 24-hour period

Age	Dose
Adults & children 12 years and older	20 mL every 4 hours
Children under 12 years of age	Do not use

**Other information**

- each 20 mL contains: sodium 10 mg
- dosage cup provided
- store between 15-30°C (59-86°F)
- do not refrigerate

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, Xanthan gum

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

\*This product is not manufactured or distributed by Reckitt Benckiser Inc., distributor of Mucinex® FAST-MAX™ Cold, Flu & Sore Throat.

**MADE WITH PRIDE & CARE FOR H-E-B®**  
SAN ANTONIO, TX 78204

**MAX COLD, FLU, AND SORE THROAT**

acetaminophen,dextromethorphan hydrobromide, guaifenesin , phenylephrine hcl liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-316
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL

**PHENYLEPHRINE HYDROCHLORIDE** (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)

PHENYLEPHRINE  
HYDROCHLORIDE

10 mg  
in 20 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYL GALLATE</b> (UNII: 8D45NN7V92)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-316-25	117 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/05/2014	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/05/2014	

**Labeler** - H E B (007924756)

**Registrant** - AptaPharma Inc. (790523323)

## Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(37808-316)

Revised: 12/2023

H E B