

JILGYUNGYI (20 TABLETS)- allantoin tablet
Haudongchun Co., Ltd.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Jilgyungyi (20 Tablets)

Allantoin (11%)

Vaginal skin protectant

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control right away. (1-800-222-1222)

Uses

- Moisturizes vaginal dryness**
- Relieves external itching**

Warnings

For vaginal use only

Do not use if you have (are)

- an allergic reaction (rashes, and pimples, redness, swelling, skin irritation, discomfort, burning, vaginal peeling or bleeding, pain in the abdomen or pelvic area, fluttering heartbeat and chest pain, lumbar pain, abnormal vaginal secretion, pollakisuria, abnormal dryness, excessive menstrual bleeding, odor and green colored vaginal secretion, hypotension, faint, and papulopustule, pelvic pain)**
- pregnant**
- hyper sensitive skin and skin ailment**
- scars on vagina**
- itching, rash, swelling redness, discomfort, burning, irritation vaginal peeling or bleeding**

When using this product

- do not use while having the monthly menstrual period**
- prior to absorption there is a risk the tablet may fall out of the vagina**
- often wash towels or underwear and do not wear reuse underwear again prior to washing due to risk of infections**
- try to avoid sexual relations when you have vaginal yeast infection, it can be aggravated**
- try not to use tampons and feminine cleansers**
- avoid contact with eye area**
- do not eat or swallow**
- do not use rubber products such as latex, condoms, and diaphragms together**
- Do not use other vaginal tablets while using Jil Gyung Yi**

If pregnant or breast-feeding, do not use.

Store at cool temperature and dry place with a closed container. Avoid direct sunlight

Overdose warning

- If you accidentally take the tablet orally, stop using this medication and see doctor immediately. Stop use and ask a doctor if you have**
- fever (over 36°C), pain in abdomen, odor, lumbar pain, abnormal secretion, nausea and abnormal shoulder pain.**

Dosage and Duration Information

☐ **For adults – 2 tablets at the same time**

☐ **Keep use 2 days consecutively and prefer to use once 10 days in 2 months. (after then, use it when you need)**

☐ **Do not use up to 6 hours prior to any sexual relation.**

☐ **Do not use up to 6 months after giving birth.**

☐ **Symptoms may vary depending on the age of the patient.Directions**

☐ **Use only adults and children 12 years age and over**

1) Wash hands before using this product

2) Do not remove moisture, open the package and vaginal tablets are placed into the vagina in much the same way as a tampon stand, squat, or lie on your back in a comfortable position then insert the loaded applicator into the vagina as far as it will comfortably go.

3) It is preferable to use the treatment during night time prior to bed.

4) Use of a sanitary pad after application, because there may be vaginal secretion 30min after application. If there are lots of secretion, change a new sanitary pad.

5) Wash the area around the vagina (the vulva) gently next day, bacteria in your vaginal secretion can cause itching.

Sodium Chloride, Soluble Collagen, Glucose, Magnesium Stearate, Aloe Barbadensis Leaf Extract, Cereus Grandiflorus (Cactus) Extract, Ginkgo Biloba Leaf Extract, Mentha Piperita (Peppermint) Leaf Extract, Centella Asiatica Extract, Sophora Flavescens Root Extract, Angelica Gigas Root Extract

Jilgyungyi (20 Tablets)

Distributed by
 KAOLIE LEE
 525 W California Ave APT 212 GARDEN CA 91203-2169
 Manufactured by
 HAUDONGCHUN CO., LTD.
 32 Seimdaeng-ro, Seodong-gu, Seoul, Seoul 06651, South Korea
 Question or Comments
 Call 1-82-70-8925-5702

10-1133723
 400949279


Jil Gyung-Yi



질경이는?

여성에게 좋은 다양한 성분이 함유되어 있어 친절하게 유지하도록 도와줍니다.
 민감한 피부로 이루어진 여성의 외음부를 친절하고 상세하게 해줍니다.

20정



제조번호 :
 사용기한 :



까지
 소비자 상담실 : 070-8962-5702

Drug Facts

Active Ingredient (in each tablet - 600mg) Purpose
 Allantoin (11%) — Vaginal skin protectant

Uses

- Moisturizes vaginal dryness
- Relieves external itching

Warnings

For vaginal use only

Do not use if you have (are)

- an allergic reaction (rashes, and pimples, redness, swelling, skin irritation, discomfort, burning, vaginal peeling or bleeding, pain in the abdomen or pelvic area, fluttering heartbeat and chest pain, lumbar pain, abnormal vaginal secretion, pollakiuria, abnormal dryness, excessive menstrual bleeding, odor and green colored vaginal secretion, hypotension, faint, and papulopustule, pelvic pain)
- pregnant
- hyper sensitive skin and skin ailment
- scars on vagina
- itching, rash, swelling redness, discomfort, burning, irritation vaginal peeling or bleeding

When using this product

- do not use while having the monthly menstrual period
- prior to absorption there is a risk the tablet may fall out of the vagina
- often wash towels or underwear and do not wear reuse underwear again prior to washing due to risk of infections
- try to avoid sexual relations when you have vaginal yeast infection, it can be aggravated
- try not to use tampons and feminine cleansers
- avoid contact with eye area
- do not eat or swallow
- do not use rubber products such as latex, condoms, and diaphragms together
- Do not use other vaginal tablets

If pregnant or breast-feeding, do not use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control right away. (1-800-222-1222)

Store at cool temperature and dry place with a closed container. Avoid direct sunlight

Overdose warning

- if you accidentally take the tablet orally, stop using this medication and see doctor immediately.

Drug Facts (Continued)

Stop use and ask a doctor if you have

- fever (over 36°C), pain in abdomen, odor, lumbar pain, abnormal secretion, nausea and abnormal shoulder pain.

Cautions

- if using the product for consecutive days, causing a temporary vaginal discharge and itching. For such case, please stop using the product and discuss with physician for continued uses. (recommended to treat spouse together) However, if you have itchy based on chronic vaginosis, please desire not to use it.

Dosage and Duration Information

- For adults — 2 tablets at the same time
- Keep use 2 days consecutively and prefer to use once 10 days in 2 months. (after then, use it when you need)
- Do not use up to 6 hours prior to any sexual relation.
- Do not use up to 6 months after giving birth.
- Symptoms may vary depending on the age of the patient.

Directions

- Use only adults and children 12 years age and over
- 1) Wash hands before using this product
- 2) Do not remove moisture, open the package and vaginal tablets are placed into the vagina in much the same way as a tampon stand, squat, or lie on your back in a comfortable position then insert the loaded applicator into the vagina as far as it will comfortably go.
- 3) It is preferable to use the treatment during night time prior to bed.
- 4) Use of a sanitary pad after application, because there may be vaginal secretion 30min after application. If there are lots of secretion, change a new sanitary pad.
- 5) Wash the area around the vagina (the vulva) gently next day, bacteria in your vaginal secretion can cause itching.

Inactive Ingredients

Sodium Chloride, Soluble Collagen, Glucose, Magnesium Stearate, Aloe Barbadensis Leaf Extract, Cereus Grandiflorus (Cactus) Extract, Ginkgo Biloba Leaf Extract, Mentha Piperita (Peppermint) Leaf Extract, Centella Asiatica Extract, Sophora Flavescens Root Extract, Angelica Gigas Root Extract



JILGYUNG YI (20 TABLETS)

allantoin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:556 18-2004
Route of Administration	VAGINAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength		
ALLANTOIN (UNII: 344S277G0Z) (ALLANTOIN - UNII:344S277G0Z)	ALLANTOIN	0.11 in 600 mg		
Inactive Ingredients				
Ingredient Name	Strength			
Sodium Chloride (UNII: 451W47IQ8X)				
.ALPHA.-D-GLUCOPYRANOSE (UNII: 5J5I9EB41E)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
ALOE VERA LEAF (UNII: ZY81Z83H0X)				
GINKGO (UNII: 19FUJ2C58T)				
MENTHA PIPERITA LEAF (UNII: A389O33LX6)				
CENTELLA ASIATICA (UNII: 7M867G6T1U)				
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)				
ANGELICA GIGAS ROOT (UNII: 32766B2FHX)				
Product Characteristics				
Color	white	Score	no score	
Shape	capsule	Size	14mm	
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55618-2004-3	2 in 1 PACKAGE	07/07/2016	
1	NDC:55618-2004-2	10 in 1 BLISTER PACK		
1	NDC:55618-2004-1	600 mg in 1 CAPSULE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved drug other		07/07/2016		

Labeler - Haudongchun Co., Ltd. (688479464)

Registrant - Haudongchun Co., Ltd. (688479464)

Establishment

Name	Address	ID/FEI	Business Operations
Haudongchun Co., Ltd.		688479464	manufacture(55618-2004)