CVS CHILDRENS NIGHTTIME MULTI SYMPTOM COLD- acetaminophen, diphenhydramine hydrochloride, phenylephrine hydrochloride liquid CVS PHARMACY

CVS Children's Nighttime Multi-Symptoms Cold 4 FL OZ (180 mL)

Drug Facts

Active ingredients (in each 10 mL)	Purposes	
Acetaminophen 325 mg	Pain reliever/fever reducer	
Diphenhydramine HCl 12.5 mg	Antihistamine/cough suppressant	
Phenylephrine HCl 5 mg	Nasal decongestant	

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver Warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- To make child sleepy
- with any other drug containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the child have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if the child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning:

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 years to 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in 24-hours period
- children under 6 years of age: do not use

Other information

- each 10 mL contains: sodium 4 mg
- store at room temperature
- do not refrigerate

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, natural and artificial flavor, potassium citrate, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum.

Questions or comments?

1-866-467-2748

PRINCIPAL DISPLAY PANEL

Compare to the active ingredients in Children's Mucinex ${}^{\circledR}$ Night Time Multi-Symptom Cold

NDC# 51316-733-04

Children's

Night TimeMuti-Symptom Cold

AcetaminophenPain Reliever/Fever Reducer
 Diphenhydramine HCl Antihistamine/Cough Suppressant
 Phenylephrine HCl Nasal Decongestant

Relieves Stuffy Nose

Controls Cough

Relieves Fever & Sore Throat

Relieves Runny Nose & Sneezing

Dosing Cup Included

For Ages 6 & Over

Very Berry Flavor

Naturally and Artificially Flavored

4 FL OZ (118 mL)

*This product is not manufactured or distributed by RB Health (US), distributor Children's Mucinex [®]Nighttime Multi-Symptoms Cold..

TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

Distributed by:



acetaminophen, diphenhydramine hydrochloride, phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-733
--------------	----------------	--------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg in 10 mL	
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 10 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
EDETATE DISODIUM (UNII: 7FLD91C86K)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
POTASSIUM CITRATE (UNII: EE90ONI6FF)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
PROPYL GALLATE (UNII: 8D4SNN7V92)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:51316-733- 04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/20/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M010	02/20/2023	

Labeler - CVS PHARMACY (062312574)

Revised: 12/2025 CVS PHARMACY