

**CITROMA- magnesium citrate liquid**  
**Old East Main Co.**

-----  
**Rexall 164.000/164AA Rev 2 - AB**  
**Cherry Magnesium Citrate**

**Active ingredient (in each fl oz)**

Magnesium citrate 1.745 g

**Purpose**

Saline laxative

**Uses**

- for relief of occasional constipation (irregularity).
- Generally produces bowel movement in 1/2 to 6 hours

**Warnings**

for this product

**Ask a doctor before use if you have**

- kidney disease
- a magnesium restricted diet
- abdominal pain, nausea, or vomiting
- noticed a sudden change in bowel habits that persists over a period of 2 weeks
- already used a laxative for a period longer than 1 week

**Ask a doctor or pharmacist before use if you are**

- taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if**

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

**if pregnant or breast-feeding,**

ask help professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Storage**

Store at temperatures between 46° and 86°F (8° and 30° C)

**Directions**

- shake well before using
- drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age - 2 to 3 fl oz maximum 3 fl oz in 24 hours

children under 2 years of age - ask a doctor

**other information**

- **each fl oz contains:** magnesium 290 mg
- **each fl oz contains:** sodium 1 mg

**Inactive ingredients**

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

**Questions?**

Call 1-866-4-REXALL

**ADVERSE REACTION**

Non-USP: Non-Carbonated, Non-Pasteurized; Contains Preservatives

Since 1903 Rexall<sup>®</sup> is a registered trademark of Rexall Sundown, Inc.

Visit us at: [Rexall.com](http://Rexall.com) or call 1-866-4-REXALL

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

**Principal display panel**

Since 1903

Rexall®

Magnesium

Citrate

Saline Laxative

- Oral Solution
- Relief of occasional constipation

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECK RING

Cherry Flavored

10 FL OZ (296 mL)

**Since 1903 Rexall®**  
**Magnesium Citrate Saline Laxative**  
Cherry Flavored  
10 FL OZ (296 mL)

• Oral solution  
• Relief of occasional constipation

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN OR SEPARATED FROM NECK RING

| Drug Facts   | Drug Facts (continued)   |
|--|--|
| <b>Active ingredient (in each fl oz) Purpose</b><br>Magnesium citrate 1.745 g Saline laxative  | or contact a Poison Control Center right away.<br>Store at temperatures between 48° and 88° F (8° and 30° C)   |
| <b>Uses</b> ■ for relief of occasional constipation (irregularity).<br>■ Generally produces bowel movement in 1/2 to 6 hours.  | <b>Directions</b> ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses   |
| <b>Warnings</b><br>Ask a doctor before use if you have:<br>■ kidney disease ■ a magnesium restricted diet<br>■ abdominal pain, nausea, or vomiting<br>■ noticed a sudden change in bowel habits that persists over a period of 2 weeks<br>■ already used a laxative for a period longer than 1 week<br>Ask a doctor or pharmacist before use if you are:<br>■ taking any other drug.<br>■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.<br>Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.<br>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help. | adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours<br>children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours<br>children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours<br>children under 2 years of age ask a doctor |
|  | <b>Other information</b><br>■ each fl oz contains: magnesium 290 mg<br>■ each fl oz contains: sodium 1 mg  |
|  | <b>Inactive ingredients</b> benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water   |
|  | <b>Questions? Call 1-866-4-REXALL</b>  |

Visit us at: [Rexall.com](http://Rexall.com) or call 1-866-4-REXALL  
DISTRIBUTED BY OLD EAST MAIN CO. 100 MISSION RIDGE GOODLETTSVILLE, TN 37072 L0017580SD

## CITROMA

magnesium citrate liquid

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:55910-961 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength           |
|--|-------------------|--------------------|
| <b>MAGNESIUM CITRATE</b> (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838) | MAGNESIUM CITRATE | 1.745 g in 29.6 mL |

### Inactive Ingredients

| Ingredient Name                                      | Strength |
|--|----------|
| <b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)               |          |
| <b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)    |          |
| <b>EDETATE DISODIUM ANHYDROUS</b> (UNII: 8NLQ36F6MM) |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                  |          |
| <b>WATER</b> (UNII: 059QF0KO0R)                      |          |

### Packaging

| # | Item Code        | Package Description  | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:55910-961-38 | 296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 02/18/2018           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M007                                     | 02/18/2018           |                    |

**Labeler** - Old East Main Co. (068331990)

**Registrant** - Nice-Pak Products, LLC (119091520)

### Establishment

| Name                   | Address | ID/FEI    | Business Operations    |
|------------------------|---------|-----------|------------------------|
| Nice-Pak Products, LLC |         | 119091514 | manufacture(55910-961) |

Revised: 3/2026

Old East Main Co.