

FOAMING HAND WASH- chlorhexidine gluconate liquid
Medline Industries, LP

994 CHG Foaming Hand Wash

Active ingredient

Chlorhexidine gluconate 2% solution

Purpose

Antiseptic

Uses

- • **surgical hand scrub**: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- • **healthcare personnel handwash**: helps reduce bacteria that potentially can cause disease

Warnings

For external use only

Allergy alert: This product may cause a severe allergic reaction. Symptoms may include:

- • wheezing/difficulty breathing
- • shock
- • facial swelling
- • hives
- • rash

If an allergic reaction occurs, stop use and seek medical help right away

Do not use

if you are allergic to chlorhexidine gluconate or any other ingredient in this product

When using this product

- • keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- • if solution should contact these areas, rinse out promptly and thoroughly with water
- • do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- • wet hands and forearms with water
- • scrub for 3 minutes with 5 mL of product (7 strokes) and a wet brush, paying close attention to the nails, cuticles and skin between the fingers
- • a separate nail cleaner may be used
- • rinse thoroughly
- • wash for an additional 3 minutes with 5 mL of product (7 strokes) and rinse under running water and dry thoroughly

Healthcare personnel handwash:

- • wet hands with water
- • dispense 5 mL of product (7 strokes) into cupped hands and wash in a vigorous manner for 15 seconds
- • rinse and dry thoroughly

Other information

- • store at 20-25°C (68-77°F)
- • avoid excessive heat above 40°C (104°F)

Inactive ingredients

citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

Manufacturing Information

Manufactured for: Medline Industries, LP

Three Lakes Drive, Northfield, IL 60093 USA

Made in USA

www.medline.com

1-800-MEDLINE

REF: HHCHG1000FM

V2 RA22XLI

Package Label

NDC: 53329-994-84



SPECTRUM™

Foaming Hand Wash

CLINICAL



Chlorhexidine Gluconate
2% Solution - Antiseptic

Manufactured for Medline Industries, LP, Three Lakes Drive, Northfield, IL
60093 USA. www.medline.com V2 RA22XLI



1 93489 10251 5

See Front for Instructions for Use and *Drug Facts*

33.8 fl oz
(1.05 qt) (1 Liter)

L0015924FB

2MED1LBBTLLBL



Foam Hand Wash

NDC: 53329-994-84

CHLORHEXIDINE GLUCONATE
2% SOLUTION - ANTISEPTIC with Foam Pump

Helps Reduce Bacteria



Drug Facts

Active ingredient

Chlorhexidine gluconate 2% solution Antiseptic

Purpose

Uses • surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
• healthcare personnel handwash: helps reduce bacteria that potentially can cause disease

Warnings

For external use only

Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing

- shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away

Do not use if you are allergic to chlorhexidine gluconate or any other ingredient in this product

When using this product • keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.

- if solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub: • wet hands and forearms with water • scrub for 3 minutes with 5 mL of product (7 strokes) and a wet brush, paying close attention to the nails, cuticles and skin between the fingers • a separate nail cleaner may be used • rinse thoroughly

• wash for an additional 3 minutes with 5 mL of product (7 strokes) and rinse under running water and dry thoroughly

Healthcare personnel handwash: • wet hands with water • dispense 5 mL of product (7 strokes) into cupped hands and wash in a vigorous manner for 15 seconds • rinse and dry thoroughly

Other information • store at 20-25°C (68-77°F) • avoid excessive heat above 40°C (104°F)

Inactive ingredients citric acid, cocamide DEA, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water

Net Contents: 33.8 fl oz (1.05 qt) (1 Liter)

HHCHG1000FM

V2 RA22XLI

2MED1LFBTLBL

FOAMING HAND WASH

chlorhexidine gluconate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53329-994
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
COCO DIETHANOLAMIDE (UNII: 92005F972D)	
LAURAMINE OXIDE (UNII: 4F6FC4MI8W)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
HYDROXYETHYL CELLULOSE (2000 CPS AT 1%) (UNII: S38J6RZN16)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53329-994-84	1000 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019422	01/01/2020	

Labeler - Medline Industries, LP (025460908)**Registrant** - Medline Industries, LP (025460908)

Revised: 2/2022

Medline Industries, LP