

**COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid**  
**WinCo Foods, LLC**

-----

**Drug Facts**

**Active ingredients (in each 15 mL)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

if a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- a sodium-restricted diet
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if**

you are taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 Days (adult)
- fever gets worse, or last more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs, with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center

(1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter

---

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to 6 years	ask a doctor
children under 4 years	do not use

---

- **When using Day Time and Night Time products, carefully read each label to ensure correct dosing**

### Other information

- **each 15 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

### Inactive ingredients

citric acid, FD&C yellow 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

### Questions?

Call **800-824-1706** Monday-Friday 9am-4pm MST

### Principal Display Panel

\*Compare to the active ingredients in VICKS® DAYQUIL® Cold & Flu

### Multi-Symptom

### Cold & Flu Relief

### Daytime

ACETAMINOPHEN 325 mg /

PAIN RELIEVER- FEVER REDUCER

DEXTROMETHORPHAN HBr 10 mg/

COUGH SUPPRESSANT

PHENYLEPHRINE HCl 5 mg/

NASAL DECONGESTANT

For Ages 6 Years & Over

Alcohol Free

Antihistamine-free

Daytime

Original Flavor

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY:

WINCO FOODS, LLC,

BOISE, ID 83704

**Package Label**

\*Compare to the active ingredients in  
VICKS® DAYQUIL® Cold & Flu

**WinCo**  
FOODS

MULTI-SYMPTOM  
**Cold & Flu  
Relief**

**ACETAMINOPHEN 325 mg/**

PAIN RELIEVER - FEVER REDUCER

DEXTROMETHORPHAN HBr 10mg/

COUGH SUPPRESSANT

PHENYLEPHRINE HCl 5 mg /

NASAL DECONGESTANT

For Ages 6 Years & Over

Alcohol Free

Antihistamine-free

**Daytime**

Original Flavor

12 FL OZ (355 mL)

P-D-A380D LB00863

**Drug Facts**

**Active ingredients  
(in each 15 mL)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Purposes**

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

**PARENTS:**

Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

DISTRIBUTED BY:  
WINCO FOODS, LLC,  
BOISE, ID 83704



**100%  
SATISFACTION  
GUARANTEE**  
Comments & Questions?  
[www.wincofoods.com](http://www.wincofoods.com)

WF2408 PLD-A380D LB010068



**PEEL CORNER FOR MORE DRUG FACTS**

**Drug Facts (continued)**

**Uses**

- temporarily relieves common cold and flu symptoms
- minor aches and pains
- headache
- sore throat
- nasal congestion
- fever
- cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours

taken with other drugs containing acetaminophen

adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**PEEL CORNER FOR MORE DRUG FACTS**

**Drug Facts (continued)**

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Drug Facts (continued)**

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

**when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

**Drug Facts (continued)**

**Other information**

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions?**

Call 800-824-1706 Monday - Friday 9am - 4pm MST

## WINCO Multi Symptom Cold and Flu Relief

### COLD AND FLU RELIEF MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-287
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

#### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-287-54	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/28/2017	

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/28/2017	

---

**Labeler** - WinCo Foods, LLC (056098817)

Revised: 9/2024

WinCo Foods, LLC