TUSSIN DMMAXIMUM STRENGTH - dextromethorphan hbr, guaifenes in liquid CVS Pharmacy

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Maximum Strength Tussin DM

Active ingredients (in each 5 mL tsp)

Dextromethorphan HBr, USP 10mg Guaifenesin, USP 200mg

Purposes

Cough suppressant

Expectorant

Uses

helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes temporarily relieves:

- cough due to minor bronchial irritation as may occur with a cold or inhaled irritants
- the impulse to cough

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions or parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if

cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- shake well before using
- do not take more than 6 doses in any 24-hour period
- only use the measuring cup that is included with this product for accurate dosing

Age	Dose
adults and children 12	
years and over	every 4 hours
children under 12	do notuco
years	do not use

Other information

- each teaspoon contains: sodium 5 mg
- store at 20-25°C (68-77°F)
- do not refrigerate
- alcohol-free
- measuring cup provided

Inactive ingredients:

artificial and natural flavors, carboxymethylcellulose sodium, citric acid, Dand C red no. 33, FD and c red no. 40, glycerin, high fructose corn syrup, menthol, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, xanthan gum

Principal Display panel

NDC 59779-967-04

Compare to the active ingredients in Adult Robitussin Cough and Chest congestion

Maximum Strength
Tussin DM
Adult Cough and chest congestion

Cough Supressant (Dextromethorphan HBr) Expectorant (Guaifenesin)

Relieves: Cough Chest congestion/mucus alcohol free

Non-Drowsy



1 image



DO NOT USE IF SEAL UNDER CAP IMPRINTED WITH
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each 5 mL tsp) Purposes
Dextromethorphan HBr, USP 10 mg Cough suppressant
Guaifenesin, USP 200 mg Expectorant

Uses

- temporarily relieves
 - cough due to minor bronchial irritation as may occur with a cold or inhaled irritants
 - the impulse to cough
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- shake well before using
- do not take more than 6 doses in any 24-hour period
- only use the measuring cup that is included with this product for accurate dosing

Age	Dose	
adults & children 12 years & over	2 tsp (10 mL) every 4 hours	
children under 12 years	do not use	

Other Information

- each teaspoon contains; sodium 5 mg
- store at 20-25°C (68-77°F)
- do not refrigerate
- alcohol-free
- measuring cup provided

Inactive ingredients artificial & natural flavors, carboxymethylcellulose sodium, citric acid, D&C red no. 33, FD&C red no. 40, glycerin, high fructose corn syrup, menthol, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, xanthan gum.

*This product is not manufactured or distributed by Wyeth Consumer Healthcare, the distributor of Adult ROBITUSSIN® Maximum Strength Cough + Chest Congestion DM.

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2013 CVS/pharmacy www.cvs.com 1-800-shop-CVS

V-11496







2 image

TUSSIN DMMAXIMUM STRENGTH

dextromethorphan hbr, guaifenesin liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:59779-967

Route of Administration

ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL		
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 5 mL		

Inactive Ingredients				
Ingredient Name	Strength			
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679 OBS 311)				
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)				
D&C RED NO. 33 (UNII: 9DBA0SBB0L)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
HIGH FRUCTOSE CORN SYRUP (UNII: XY6 UN3QB6S)				
MENTHOL (UNII: L7T10EIP3A)				
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
XANTHAN GUM (UNII: TTV12P4NEE)				

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:59779-967-03	1 in 1 CARTON			
1		118 mL in 1 BOTTLE			
2	NDC:59779-967-04	1 in 1 CARTON			
2		237 mL in 1 BOTTLE			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	07/01/2009			

Labeler - CVS Pharmacy (062312574)

Revised: 6/2013 CVS Pharmacy