

**GOOD SENSE ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated  
L. Perrigo Company**

-----  
**Perrigo Anti-Diarrheal Drug Facts**

**Active ingredient (in each caplet)**

Loperamide HCl 2 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

|  |  |
|--|--|
| adults and children<br>12 years and over | 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours  |
| children 9-11 years<br>(60-95 lbs)       | 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours |
| children 6-8 years<br>(48-59 lbs)        | 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours |
| children 2-5 years<br>(34 to 47 lbs)     | ask a doctor   |
| children under 2 years<br>(up to 33 lbs) | do not use   |

**Other information**

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

**Inactive ingredients**

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

Anti-Diarrheal

See New Warnings

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls the Symptoms of Diarrhea

Actual Size

Compare to active ingredient of Imodium<sup>®</sup> A-D

100% SATISFACTION GUARANTEED

12 Caplets\*

\*Capsule-Shaped Tablets

GOODSENSE/ANTI-DIARRHEAL

22453 C2 C8

OPEN OTHER END

**Drug Facts (continued)**

**Inactive ingredients** anhydrous lactose, carnauba wax, D & C yellow no. 10 aluminum lake, FD & C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Gluten Free

**Drug Facts (continued)**

**Questions or comments?** 1-800-71-9-260

**Drug Facts**

**Active ingredient** (in each caplet) Loperamide HCl 2 mg.....Anti-diarrheal

**Purpose** Use to control symptoms of diarrhea, including Traveler's Diarrhea

**Warnings** Allergic alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. **Heart alert:** Taking more than directed can cause serious heart problems or death. Do not use if you have had body or back stool. Ask a doctor before use if you have fever, mucus in the stool, a history of liver disease, a history of abnormal heart rhythm. Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** dizziness, drowsiness or lightheadedness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if** symptoms get worse, diarrhea lasts for more than 2 days, you get abdominal swelling or bloating. These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

**Other information**

- Store at 20°-25°C (68°-77°F)
- See end panel for lot number and expiration date

**Directions**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

- Find right dose on chart, if possible, use weight to dose; otherwise, use age.

|                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours  |
| children 9-11 years (60-95 lbs)       | 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours |
| children 6-8 years (48-59 lbs)        | 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours |
| children 2-5 years (34 to 47 lbs)     | ask a doctor   |
| children under 2 years (up to 33 lbs) | do not use   |



DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

Distributed By **Perrigo** Allegan, MI 49010

GOODSENSE®

Loperamide Hydrochloride Tablets, 2 mg

# Anti-Diarrheal

GOODSENSE®


NDC 0113-0224-53

Anti-Diarrheal

See New Warnings

Loperamide Hydrochloride  
Tablets, 2 mg





## GOOD SENSE ANTI DIARRHEAL

loperamide hydrochloride tablet, film coated

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:0113-0224 |
| <b>Route of Administration</b> | ORAL           |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength        | Strength |
|---|--------------------------|----------|
| <b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II) | LOPERAMIDE HYDROCHLORIDE | 2 mg     |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)                |          |
| <b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)                     |          |
| <b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)        |          |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)               |          |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)       |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A) |          |
| <b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)            |          |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)              |          |

### Product Characteristics

|                 |       |                     |          |
|-----------------|-------|---------------------|----------|
| <b>Color</b>    | GREEN | <b>Score</b>        | 2 pieces |
| <b>Shape</b>    | OVAL  | <b>Size</b>         | 10mm     |
| <b>Flavor</b>   |       | <b>Imprint Code</b> | L2       |
| <b>Contains</b> |       |                     |          |

**Packaging**

| # | Item Code        | Package Description                                    | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:0113-0224-53 | 12 in 1 CARTON   | 03/08/2003           |                    |
| 1 |                  | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |
| 2 | NDC:0113-0224-62 | 24 in 1 CARTON   | 03/25/2010           |                    |
| 2 |                  | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |
| 3 | NDC:0113-0224-89 | 18 in 1 CARTON   | 03/06/2003           | 03/05/2019         |
| 3 |                  | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |
| 4 | NDC:0113-0224-91 | 6 in 1 CARTON  | 02/25/2003           |                    |
| 4 |                  | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

**Marketing Information**

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA075232                               | 02/25/2003           |                    |

**Labeler** - L. Perrigo Company (006013346)

Revised: 11/2022

L. Perrigo Company