

ASPIRIN 81 MG- aspirin tablet, coated
Beacon Pharma, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Aspirin 81 mg Tablets

Active ingredient
(in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease

- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

(609)375-2548

Principal Display Panel

NDC 72629-020-10, 100 Count

<p>NDC 72629-020-10</p> <p>BEACON DISTRIBUTION</p> <p>Generic for Bayer Aspirin 81mg</p> <p>Low Dose</p> <h1>Aspirin</h1> <p>Pain Reliever / Fever Reducer(NSAID) Fast, Safe Pain Relief</p> <p>100 ENTERIC-COATED TABLETS</p>	<p>Drug Facts</p> <p>Active Ingredient (in each tablet) Purpose Aspirin 81 mg (NSAID).....Pain Reliever *nonsteroidal anti-inflammatory drug</p> <p>Uses • temporarily relieves minor aches and pains • for other uses, see your doctor but do not use for more than 10 days without consulting your doctor because serious side effects may occur</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: • hives • facial swelling • shock • asthma (wheezing) Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed</p>	<p>Drug Facts (continued)</p> <p>• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)</p> <p>Do not use • if you are allergic to aspirin or any other pain reliever/fever reducer</p> <p>Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have not been drinking fluids • you have lost a lot of fluid due to vomiting or diarrhea</p> <p>Ask a doctor or a pharmacist before use if you are • taking a prescription drug for diabetes, gout, or arthritis • taking any other drug • under a doctor's care for any serious condition</p> <p>Stop use and ask a doctor if • you experiencing any of the following signs of stomach bleeding: • feel faint • have bloody or black stools • vomit blood • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area • any new symptoms appear • ringing in the ears if a loss of hearing occurs</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children: In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions • drink a full glass of water with each dose • adults and children 12 years and over; take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor • children under 12 years; consult a doctor</p> <p>Other information • store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°) • use by expiration date on package</p> <p>Inactive Ingredients anhydrous lactose, black iron oxide, brown iron oxide, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, hypromellose, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polyorbate 80, potassium hydroxide, propylene glycol, shellac, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate</p>	<p>Distributed by: Beacon Distribution LLC Pennington NJ 08534</p> <p>TEMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING</p> <p>8 60142 1002 12 2</p> <p>Lot No.: T-18187 Exp. Date: 02/2022</p>
--	--	--	---	---

NDC 72629-020-30, 300 Count

<p>NDC 72629-020-30</p> <p>BEACON DISTRIBUTION</p> <p>Generic for Bayer Aspirin 81mg</p> <p>Low Dose</p> <h1>Aspirin</h1> <p>Pain Reliever / Fever Reducer(NSAID) Fast, Safe Pain Relief</p> <p>300 ENTERIC-COATED TABLETS</p>	<p>Drug Facts</p> <p>Active Ingredient (in each tablet) Purpose Aspirin 81 mg (NSAID).....Pain Reliever *nonsteroidal anti-inflammatory drug</p> <p>Uses • temporarily relieves minor aches and pains • for other uses, see your doctor but do not use for more than 10 days without consulting your doctor because serious side effects may occur</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: • hives • facial swelling • shock • asthma (wheezing) Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed</p>	<p>Drug Facts (continued)</p> <p>• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)</p> <p>Do not use • if you are allergic to aspirin or any other pain reliever/fever reducer</p> <p>Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have not been drinking fluids • you have lost a lot of fluid due to vomiting or diarrhea</p> <p>Ask a doctor or a pharmacist before use if you are • taking a prescription drug for diabetes, gout, or arthritis • taking any other drug • under a doctor's care for any serious condition</p> <p>Stop use and ask a doctor if • you experiencing any of the following signs of stomach bleeding: • feel faint • have bloody or black stools • vomit blood • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area • any new symptoms appear • ringing in the ears if a loss of hearing occurs</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children: In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions • drink a full glass of water with each dose • adults and children 12 years and over; take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor • children under 12 years; consult a doctor</p> <p>Other information • store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°) • use by expiration date on package</p> <p>Inactive Ingredients anhydrous lactose, black iron oxide, brown iron oxide, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, hypromellose, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polyorbate 80, potassium hydroxide, propylene glycol, shellac, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate</p>	<p>Distributed by: Beacon Distribution LLC Pennington NJ 08534</p> <p>TEMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING</p> <p>8 60142 1002 10 8</p> <p>Lot No.: T-18188 Exp. Date: 02/2022</p>
--	--	--	---	---

NDC 72629-020-50, 500 Count



NDC 72629-020-50

Low Dose

Generic for Bayer Aspirin 81mg

Aspirin

Pain Reliever / Fever Reducer(NSAID)
Fast, Safe Pain Relief

500 ENTERIC-COATED TABLETS

Drug Facts

Active Ingredient (in each tablet) Aspirin 81 mg (NSAID)
Purpose Pain Reliever / Nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains for other uses, see your doctor but do not use for more than 10 days without consulting your doctor because serious side effects may occur

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
• hives • facial swelling • shock • asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Drug Facts (continued)

• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)

Do not use • if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

• stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have not been drinking fluids • you have lost a lot of fluid due to vomiting or diarrhea

Ask a doctor or a pharmacist before use if you are

• taking a prescription drug for diabetes, gout, or arthritis
• taking any other drug
• under a doctor's care for any serious condition

Stop use and ask a doctor if

• you experiencing any of the following signs of stomach bleeding: • feel faint • have bloody or black stools • vomit blood • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area • any new symptoms appear • ringing in the ears or a loss of hearing occurs

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children; In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• drink a full glass of water with each dose
• adults and children 12 years and over; take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
• children under 12 years; consult a doctor

Other information

• store at 25°C (77°F) excursions permitted between 15°-30° (59°-86°) • use by expiration date on package

Inactive Ingredients

anhydrous lactose, black iron oxide, brown iron oxide, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, hypromellose, iron oxide ochre, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, potassium hydroxide, propylene glycol, shellac, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate



8 60142 00211 5

Distributed by: Beacon Distribution LLC
Pennington, NJ, 08524

TEMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Lot No.: T-18187
Exp. Date: 02/2022

ASPIRIN 81 MG

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72629-020
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIMETHICONE (UNII: 92RU3N3Y1O)	

WATER (UNII: 059QF0KO0R)

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	S1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72629-020-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2019	
2	NDC:72629-020-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2011	
3	NDC:72629-020-30	300 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	05/01/2011	

Labeler - Beacon Pharma, Inc. (116009701)

Revised: 1/2020

Beacon Pharma, Inc.