

**LOW DOSE ASPIRIN LOW DOSE- aspirin tablet, delayed release  
SMART SENSE (KMART)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**STS - 1146 - 2019-1008**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID <sup>1</sup>)

---

<sup>1</sup> nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- temporarily relieves minor aches and pains

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours; do not exceed 48 tablets in 24 hours
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information

**Inactive ingredients**

acetylated monoglycerides, anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C red #40, FD&C yellow #6, hypromellose, hypromellose phthalate, iron oxide yellow, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polyethylene glycol, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

**PRINCIPAL DISPLAY PANEL**

Smart Sense™

Compare to St. Joseph® Low Dose Aspirin†

Low Dose Aspirin

Pain Reliever (NSAID)

Safety-Coated

180 Tablets Enteric Coated

81 mg

COATING FREE AREA

**Drug Facts**  
**Active ingredient (in each tablet)** Purpose  
 Aspirin 81 mg (NSAID).....Pain reliever  
 \*nonsteroidal anti-inflammatory drug

**Uses** ■ Temporarily relieves minor aches and pains

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product

**Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer**

**Ask a doctor before use if**  
 ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

COMPARE TO ST. JOSEPH® LOW DOSE ASPIRIN†

**smart sense**

**low dose aspirin**

PAIN RELIEVER (NSAID)  
SAFETY-COATED

**180 TABLETS**

**81 mg**

ENTERIC COATED

COMPARE TO ST. JOSEPH® LOW DOSE ASPIRIN†

**smart sense**

**low dose aspirin**

PAIN RELIEVER (NSAID)  
SAFETY-COATED

**180 TABLETS**

**81 mg**

ENTERIC COATED

**Drug Facts (continued)**  
**Inactive ingredients** acetylated monoglycerides, anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C red #40, FD&C yellow #6, hypromellose, hypromellose phthalate, iron oxide yellow, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polyethylene glycol, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

Distributed by: Knarr Corporation F114601STS\_R1  
 Hoffman Estates, IL 60179  
 Shop knarr.com

**SATISFACTION GUARANTEE**  
 If you are unsatisfied for any reason, return the unused portion to the store for a full refund or call 1-800-842-7886.

INK AND COATING FREE FOR LOT AND EXPIRATION STAMPING

8 83967 38784 7

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

†This product is not manufactured or distributed by St. Joseph Health Products, LLC, distributor of St. Joseph® Low Dose Aspirin.

**Drug Facts (continued)**  
 ■ you are taking a diuretic ■ you have asthma  
**Ask a doctor or pharmacist before use if you are** taking a prescription drug for:  
 ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
 ■ an allergic reaction occurs. Seek medical help right away.  
 ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ pain gets worse or lasts for more than 10 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.  
**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours; do not exceed 48 tablets in 24 hours ■ children under 12 years: do not use unless directed by a doctor

**Other information**  
 ■ Store between 20°-25°C (68°-77°F) in a dry place ■ retain carton for complete product information

**LOW DOSE ASPIRIN LOW DOSE**  
 aspirin tablet, delayed release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49738-786
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
DIACETYLATED MONO GLYCERIDES (UNII: 5Z17386USF)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
HYPROMELLOSE PHTHALATE (24% PHTHALATE, 55 CST) (UNII: 87Y6436BKR)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

**Product Characteristics**

Color	orange (PEACH)	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	heart
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49738-786-18	1 in 1 CARTON	02/01/2013	
1		180 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:49738-786-36	2 in 1 CARTON	07/01/2011	07/31/2019
2		180 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	06/30/2011	

**Labeler** - SMART SENSE (KMART) (008965873)

Revised: 10/2019

SMART SENSE (KMART)