

ANTI-DIARRHEAL- loperamide hydrochloride solution

Safeway

Better Living Brands Anti-Diarrheal Oral Solution LLC Drug Facts

Active ingredient (in each 7.5 mL)

Loperamide HCl 1 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Signature SELECT™

COMPARE TO Imodium® A-D active ingredient

LOPERAMIDE HYDROCHLORIDE ORAL SOLUTION 1mg per 7.5 mL

ANTI-DIARRHEAL ORAL SOLUTION

MINT FLAVOR

Controls the symptoms of diarrhea

Anti-Diarrheal Oral Solution

LOVE IT OR IT'S ON US

SEE NEW WARNING AND DIRECTIONS

8 FL OZ (240 mL)

Drug Facts

Active ingredient Purpose (in each 7.5 mL)

Loperamide HCl 1 mg.....Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Peel open to read complete directions and warnings before purchase

NDC 21130-381-34

Signature
SELECT™

LOPERAMIDE HYDROCHLORIDE
ORAL SOLUTION 1 mg per 7.5 mL

ANTI-DIARRHEAL ORAL SOLUTION

MINT FLAVOR

Controls the symptoms
of diarrhea
Anti-Diarrheal Oral Solution

SEE NEW WARNING
AND DIRECTIONS

8 FL OZ (240 mL)

COMPARE TO
Imodium® A-D
active ingredient*

Do not use if printed
plastic neckband is
broken or missing.

*This product is not
manufactured or distributed
by Johnson & Johnson
Consumer Inc., distributor of
Imodium® A-D.

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
†1-888-723-3929

LOVE IT OR IT'S ON US!



Scan here for more
information

S 1872 RD 24044

EXP LOT

: 64534 2A F1



3

Drug Facts (continued)

Ask a doctor before use if you have

- fever ■ mucus in the stool ■ a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
 - diarrhea lasts for more than 2 days
 - you get abdominal swelling or bulging.
- These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children
12 years and over

30 mL after the first loose stool;
15 mL after each subsequent
loose stool; but no more than
60 mL in 24 hours

children 9-11 years
(60-95 lbs)

15 mL after the first loose stool;
7.5 mL after each subsequent
loose stool; but no more than
45 mL in 24 hours

Drug Facts (continued)

children 6-8 years
(48-59 lbs)

15 mL after the first loose stool;
7.5 mL after each subsequent
loose stool; but no more than
30 mL in 24 hours

children 2-5 years
(34 to 47 lbs)

ask a doctor

children under
2 years
(up to 33 lbs)

do not use

Other information

- each 30 mL contains: sodium 15 mg
- store between 20-25°C (68-77°F)

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-800-719-9260

ANTI-DIARRHEAL				
loperamide hydrochloride solution				
Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:21130-381
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)			LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL
Inactive Ingredients				
Ingredient Name				Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
GLYCERIN (UNII: PDC6A3C0OX)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
XANTHAN GUM (UNII: TTV12P4NEE)				
DIMETHICONE (UNII: 92RU3N3Y1O)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
Product Characteristics				
Color	GREEN (opaque, viscous)		Score	
Shape			Size	
Flavor	MINT		Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-381-34	240 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/15/2024	
Marketing Information				
Marketing		Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
ANDA	ANDA091292	10/15/2024	

Labeler - Safeway (009137209)

Revised: 10/2024

Safeway