STOOL SOFTENER DOCUSATE SODIUM- docusate sodium capsule, gelatin coated

Magno-Humphries, Inc.

Stool Softener Docusate Sodium

Drug Facts

Active ingredient (in each softgel)

Docusate sodium 100 mg

Purpose

Stool softener

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 12 to 72 hours

Warnings

Do not use

• if you are presently taking mineral oil, unless told to do so by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over 2 weeks

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a stool softener laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

• Take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses.

adults and children 12 years of age and older	take 1 to 3 softgels daily
children 2 to under 12 years of age	take 1 softgel daily
children under 2 years	ask a doctor

Other information

- ach softgel contains: sodium 7mg
- store at 15° to 30°C (59° to 86°F)
- protect from moisture

Inactive ingredients

FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, purified water, sorbitol special

Questions?

call toll-free 1-800-935-6737

Package Labeling:

NDC 54257-902-02 SCOFFICENCE SOFFICENCE Compare to the active ingredient in Colace®* Effective, Gentle, Stimulant Free 100 SOFTGELS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each softgel) Purpose
Docusate sodium 100 mg......Stool softener

Uses ■ relieves occasional constipation (irregularity) ■ generally produces bowel movement in 12 to 72 hours

Warnings

Do not use ■ if you are presently taking mineral oil, unless told to do so by a doctor

Ask a doctor before use if you have: ■ stomach pain ■ nausea ■ vomiting ■ noticed a sudden change in bowel habits that lasts over 2 weeks

Stop use and ask a doctor if ■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition. ■ you need to use a stool softener laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ Take only by mouth. May be taken as a single daily dose or in divided doses.

of age and older	take 1 to 3 softgels daily	
children 2 to under 12 years of age	take 1 softgel daily	
children under 2 years	ask a doctor	

Other information ■ each softgel contains: sodium 7mg ■ store at 15° to 30°C (59° to 86°F)

protect from moisture

Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, purified water, sorbitol special

Questions? call toll-free 1-800-935-6737

*This product is not manufactured or distributed by the owner of the registered trademark Colace_®



DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each softgel) Purpose

Docusate sodium 100 mg.....Stool softener

Uses ■ relieves occasional constipation (irregularity) ■ generally produces bowel movement in 12 to 72 hours

Warnings

Do not use ■ if you are presently taking mineral oil, unless told to do so by a doctor

Ask a doctor before use if you have: ■ stomach pain ■ nausea ■ vomiting ■ noticed a sudden change in bowel habits that lasts over 2 weeks

Stop use and ask a doctor if ■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition. ■ you need to use a stool softener laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ Take only by mouth. May be taken as a single daily dose or in divided doses.

adults and children 12 years of age and older	take 1 to 3 softgels daily
children 2 to under 12 years of age	take 1 softgel daily
children under 2 years	ask a doctor

Other information = each softgel contains: sodium

7mg store at 15° to 30°C (59° to 86°F)

protect from moisture

Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, purified water, sorbitol special

Questions? call toll-free 1-800-935-6737

*This product is not manufactured or distributed by the owner of the registered trademark Colace®



STOOL SOFTENER DOCUSATE SODIUM

docusate sodium capsule, gelatin coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54257-902
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	100 mg	

Inactive Ingredients				
Ingredient Name	Strength			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GELATIN (UNII: 2G86QN327L)				
GLYCERIN (UNII: PDC6A3C0OX)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
WATER (UNII: 059QF0KO0R)				

Product Characteristics			
Color	red	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	SCU1
Contains			

Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		NDC:54257-902- 02	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/24/2020	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M007	01/24/2020		
			Date	

Labeler - Magno-Humphries, Inc. (063251433)

Revised: 10/2023 Magno-Humphries, Inc.