

IBUPROFEN- ibuprofen tablet, coated
Harris Teeter

HTE - 1110C - 2019-1010

Drug Facts

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to

use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
 - take 1 caplet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 caplet, 2 caplets may be used
 - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F)
- read all warnings and directions before use
- retain carton for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, corn starch, iron oxide red, polyethylene glycol, polyvinyl alcohol, povidone K30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

Harris Teeter

Ibuprofen

Compare to the Active Ingredient in Advil(R)

Ibuprofen Tablets, 200 mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

24 CAPLETS**

**Capsule-shaped tablets

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil®

DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) Purpose
Ibuprofen 200 mg (NSAID)*...Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ toothache ■ headache
■ menstrual cramps ■ the common cold
■ muscular aches ■ minor pain of arthritis
■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives
■ facial swelling ■ skin reddening
■ asthma (wheezing) ■ blisters
■ shock

If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warnings: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers or fever reducers

■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ taking any other drug

When using this product
■ take with food or milk if stomach upset occurs
■ stop use and ask a doctor if
■ side effects occur. You may report side effects to FDA at 1-800-FDA-1088.

■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing

Drug Facts (continued)

■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over:
■ take 1 caplet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Other information
■ store at 20°-25°C (68°-77°F)
■ avoid excessive heat above 40°C (104°F)
■ read all warnings and directions before use
■ retain carton for complete product information and warnings

Drug Facts (continued)
Inactive ingredients colloidal silicon dioxide, corn starch, iron oxide red, polyethylene glycol, polyvinyl alcohol, povidone K30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-844-705-4384

yourplinsurance.
PROUDLY DISTRIBUTED BY:
HARRIS TEETER, LLC, MATTHEWS, NC 28105
1-800-432-6111 or harris@teeter.com
MADE IN INDIA

Ibuprofen
Ibuprofen Tablets, 200 mg
PAIN RELIEVER / FEVER REDUCER (NSAID)
24 CAPLETS**
**capsule-shaped tablets

IBUPROFEN ibuprofen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72036-110
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	brown	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72036-110-01	1 in 1 CARTON	02/18/2013	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:72036-110-02	1 in 1 CARTON	07/23/2012	
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:72036-110-03	1 in 1 CARTON	07/23/2012	
3		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:72036-110-05	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/09/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	07/23/2012	

