STOMACH RELIEF MAXIMUM STRENGTH- bismuth subsalicylate liquid CVS Pharmacy

Drug Facts

Active ingredient (in each 15 mL dose)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- diarrhea
- traveler's' diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - o gas
 - indigestion
 - nausea
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms ahould not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of childen.

In case of overdose, get medical help or contact a Posion Control Center (1-800-222-1222) right away.

Directions

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and over:
 - 15 mL (1dose) every 1/2 or 30 mL (2 doses) every hour as needed diarrhea/traveler's diarrhea
 - 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

Other information

- each 15 mL contains: sodium 5 mg
- each 15 mL contains: salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Pepto-Bismol® Ultra active ingredient*

Ultra

STOMACH RELIEF

Bismuth Subsalicylate 525 mg PER 15 mL

Upset Stomach Reliever/Antidiarrheal

Relieves 5 Symptoms

Heartburn, Indigestion, Upset stomach, Nausea, Diarrhea

Original Flavor

SHAKE WELL

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® Ultra is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Distributed by:

CVS Pharmacy, Inc.

One CVS Drive Woonsocket, RI 02895

Package Label





Drug Facts (continued) Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) taking other salicylate products Do not use if you have an ulcer a bleeding problem bloody or black stool Ask a doctor before use if you have ■ fever ■ mucus in the stool Ask a doctor or pharmacist before use if you are taking any drug for ■ diabetes ■ gout ■ arthritis anticoagulation (thinning the blood) When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur. Stop use and ask a doctor if symptoms get worse or last more than 2 days ringing in the ears or loss of hearing occurs diarrhea lasts more than 2 days.

Drug Facts (continued)

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over: ■ 15 mL (1 dose) every ½ hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's
- diarrhea ■ 15 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

Other information

- meach 15 mL contains; sodium 5 mg
- each 15 mL contains: salicylate 206 mg
- low sodium keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

STOMACH RELIEF MAXIMUM STRENGTH

bismuth subsalicylate liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69842-229

Route of Administration ORAL

Active Ingredient/Active Moiety Basis of Ingredient Name Strength Strength

BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, BISMUTH 525 ma SALICYLIC ACID - UNII: 0414PZ4LPZ) SUBSALICYLATE in 15 mL

active Ingredients		
Ingredient Name	Strength	
BENZOIC ACID (UNII: 85KN0B0MIM)		
D&C RED NO. 22 (UNII: 1678RKX8RT)		
D&C RED NO. 28 (UNII: 767IP0Y5NH)		
GLYCERIN (UNII: PDC6A3C0OX)		
WATER (UNII: 059QF0KO0R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		

XANTHAN GUM (UNII: TTV12P4NEE)	

ı	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:69842- 229-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2020	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M008	01/31/2020		
3 17 13				

Labeler - CVS Pharmacy (062312574)

Revised: 10/2025 CVS Pharmacy