NO DRIP SEVERE NASAL MIST- oxymetazoline spray liquid CVS Health

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredient

Oxymetazoline Hydrochloride 0.05%

Purpose

Nasal Decongestant

Uses

Temporarily relieves nasal congestion due to:

- common cold
- hay fever
- upper respiroty allergies
- temporarily relieves sinus congestion and pressure
- Shrinks swollen membranes so you can breathe more freely

Warnings

Ask a Doctor before use if you have

- heart diease
- high blood pressure
- diabetes
- thyiod diease
- trouble urinating due to enlarged rostate gland.

When using this product

- Do not use more than directed
- do not use for more than three days, Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning ,stinging, sneezing or an increase in nasal discharge may occur.
- use of this container by more than one person may spread infection.

stop use and ask a doctor if

symptoms persist.

if pregnant or breast-feeding

ask a health professional before use.

Keep out of the reach of children

if swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Before using for the first time, remove the protective cap form the tip and prime the pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostrile. Fully depress rim with a firm and even stroke, and sniff deeply.
- Wipe nozzle clean after use.
- Adults and children 6 to under 12 years of age (with adult supervision):2 or 3 sprays in each nostril not more than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- Children under 6 years of age consult a doctor.

Other information

- store between 20° and 25°C (68° to 77°F)
- retain carton for future reference on full labeling

Inactive Ingredients

Benzalkonium Chloride, Camphor, Cellulose Gum, Disodium EDTA, Disodium Phosphate, Eucalyptol, Glycerin, Menthol, Microcrystalline Cellulose and Carboxymethylcellulose Sodium, PEG-6, Povidone, Propylene Glycol, Sodium Phosphate, Water (purified)

Principal Panel 0.5 FLOZ Bottle



CVS Health No Drip Severe Nasal Mist

Nasal Spray

Oxymetazoline HCL 0.05%

12 hour Nasal Spray

Net Wt 0.5 FLOZ (15ml)





SEE CARTON FOR FULL LABELING Do not use if printed seal over Cap is torn or missing

Oxymetazoline hydrochloride 0.05% Uses • See Carton.

When using this product • do not use more Warnings Ask a doctor before use if you have . heart disease . high blood pressure uninating due to an enlarged prostate gland. thyroid disease • diabetes • trouble

person may spread infection. Stop use and If pregnant or breast-feeding, ask a health professional before use. Keep out of reach than directed • do not use for more than 3 prolonged use may cause nasal congestion lof children. If swallowed, get medical help to recur or worsen. . temporary discomfor away. Directions To use: squeeze cap at such as burning, stinging, sneezing or an use of this container by more than one days. Use only as directed. Frequent or or contact a Poison Control Center right increase in nasal discharge may occur ask a doctor if symptoms persist.

If pregnant or breast-feeding, ask a health

ask a doctor if symptoms persist

professional before use. Keep out of reach

lof children. If swallowed, get medical help

away. Directions To use: squeeze cap at

or contact a Poison Control Center right

person may spread infection. Stop use and

use of this container by more than one

to recur or worsen. • temporary discomfor

such as burning, stinging, sneezing or an

increase in nasal discharge may occur

after use. • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 every 10 to 12 hours. Do not exceed 2 dose sprays in each nostril not more often than in any 24-hour period. • children under 6 counterclockwise to remove. Secure cap base on opposite sides and twist cap years of age: ask a doctor.

after use. • adults and children 6 to under 12

counterclockwise to remove. Secure cap

base on opposite sides and twist cap

years of age (with adult supervision): 2 or 3 every 10 to 12 hours. Do not exceed 2 doses

sprays in each nostril not more often than in any 24-hour period. • children under 6

years of age: ask a doctor.

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2018 CVS/pharmacy 7908068 CVS.com® 1-800-SHOP CVS V-34744

When using this product • do not use more If pregnant or breast-feeding, ask a health professional before use. Keep out of reach counterclockwise to remove. Secure cap after use. • adults and children 6 to under 1/ years of age (with adult supervision): 2 or 3 person may spread infection. Stop use and ask a doctor if symptoms persist. every 10 to 12 hours. Do not exceed 2 doses Warnings Ask a doctor before use if you urinating due to an enlarged prostate gland. Health. have • heart disease • high blood pressure than directed • do not use for more than 3 prolonged use may cause nasal congestion lof children. If swallowed, get medical help to recur or worsen. • temporary discomfor Directions To use: squeeze cap at sprays in each nostril not more often than in any 24-hour period. • children under 6 Severe SEE CARTON FOR FULL LABELING Do not use if printed seal over Cap is torn or missing such as burning, stinging, sneezing or an use of this container by more than one or contact a Poison Control Center right away. *Directions* To 1150: contacts on the control of days. Use only as directed. Frequent or increase in nasal discharge may occur thyroid disease • diabetes • trouble base on opposite sides and twist cap Oxymetazoline hydrochloride 0.05% No Drip sal Mist years of age: ask a doctor OXYMETAZOLINE HCI 0.05% Uses • See Carton Nasal decongestant 0.5 FL 0Z (15 mL)

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2018 CVS/pharmacy 7908068 CVS.com® 1-800-SHOP CVS V-34744

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2018 CVS/pharmacy 7908068 CVS.com® 1-800-SHÓP CVS V-34744



Principal display panel- Carton label





Health. SADA

DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

Drug Facts

Active ingredient

Purpose Active ingressions

Dynetocine hydrochloride 0 10%. Nasal decongestant

US&S - temporarily relieves nacid congestion due to:

• common cost • flag fever • upper respiratory
allergies • temporarily relieves sinus congestion and
pressure • shrinks reveilen nacid memoranes so you an breathe more freely

Warnings Ask a doctor before use if you have • heart disease

And a doute to either use if you have - heart disease - high bood pressure - throat disease - disabets - high bood pressure - throat disease - disabets - throate writing due to an enterped protothe gland. When severy the product - disease were the more than 5 days. One only as directed, frequent or protonyed use may cause need competion to word or acrosses. - temporary disconficial such as burning, estinging, making or an inverses in resed disodrage may occur - see of this container by more than one person may spread effection. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop use and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses a disoter

help or comunica in Polson Control Centers cight ainage.

Directions

Single colony for first time, remove the preleasing cape
from the ligh and prime pump by dispressibly pump
from the ligh and prime pump by dispressibly pump
from been all mores. It surps, such other with thurst but
these and next per between their and exposed times.

Without timing hand, more cape in in soorth if July
cognises of me with a firm, your movive and said coopy.

With a code: close after use.

- adults and children 6 its under 12 years of age juelts
auth supervisions 2 or 3 pumps in each notifiel act
more often than every 10 to 12 hours. Do not exceed
2 does in mig 24-brow print.

- children under 6 years of age; side a doctor.

Drug Facts (portious)

Other information • store between 20" to 25"C (68" to 77"9)

retain partor for future

inactive ingredients

anactive angine columns Berzalkinian O'lorde, Camptor, Celaidore Sum, Doodum EDTA, Doodum Prognant, Eurasyposi, Oyonia, Membrol, Microsystaline Celaidore and Carbonymethylori-latore Sodium PEG-6, Provione, Progriere Oyori, Sodium Progress Oyori, Sodium Progres

Incyline Gycli, Sodium hospinin, Water jourfled; Cap removed instructions. To remove cap, squeeze cap at base on opposite sides on the two Jurn impressions with fingers. While squeezing, twist cap countertricokwise to remove. Press down to use. Refer to amove on cap.

To reapply cap, turn cap on pump clockwise until it locks.

"This product is not manufactured or distributed by Bayer HealthCare LLC, the distributor of Afrin" No Drip Severe Congestion Pump Mist.

IMPORTANT: KEEP OUTER CARTON FOR FUTURE REFERENCE ON FULL LABELING

PCVS Health.

Compare to the active ingraciant in Afrin* No Drip Severe Congestion Pump Mist* Sovere

No Drip Nasal (2) Mist

OXYMETAZOLINE HCI 0.05%

Nasal decongestant 12 HOUR PUMP MIST



CVS Quality

LOT



5001014



Severe Nasal Mist

Nasal Spray

Oxymetazoline HCL 0.05%

12 hour Nasal Spray

Net Wt 0.5 FLOZ (15ml)



Health. SADA

DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

Drug Facts

Active ingredient

Purpose Active ingressions

Dynetocine hydrochloride 0 10%. Nasal decongestant

US&S - temporarily relieves nacid congestion due to:

• common cost • flag fever • upper respiratory
allergies • temporarily relieves sinus congestion and
pressure • shrinks reveilen nacid memoranes so you an breathe more freely

Warnings Ask a doctor before use if you have • heart disease

And a doute to either use if you have - heart disease - high bood pressure - throat disease - disabets - high bood pressure - throat disease - disabets - throate writing due to an enterped protothe gland. When severy the product - disease were the more than 5 days. One only as directed, frequent or protonyed use may cause need competion to word or acrosses. - temporary disconficial such as burning, estinging, making or an inverses in resed disodrage may occur - see of this container by more than one person may spread effection. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop use and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses and eath a disoter if symptoms person. Stop uses a disoter

help or comunica in Polson Control Centers cight ainage.

Directions

Single colony for first time, remove the preleasing cape
from the ligh and prime pump by dispressibly pump
from the ligh and prime pump by dispressibly pump
from been all mores. It surps, such other with thurst but
these and next per between their and exposed times.

Without timing hand, more cape in in soorth if July
cognises of me with a firm, your movive and said coopy.

With a code: close after use.

- adults and children 6 its under 12 years of age juelts
auth supervisions 2 or 3 pumps in each notifiel act
more often than every 10 to 12 hours. Do not exceed
2 does in mig 24-brow print.

- children under 6 years of age; side a doctor.

Drug Facts (portious)

Other information • store between 20" to 25"C (68" to 77"9)

retain partor for future

inactive ingredients

anactive angine columns Berzalkinian O'lorde, Camptor, Celaidore Sum, Doodum EDTA, Doodum Prognant, Eurasyposi, Oyonia, Membrol, Microsystaline Celaidore and Carbonymethylori-latore Sodium PEG-6, Provione, Progriere Oyori, Sodium Progress Oyori, Sodium Progres

Incyline Gycli, Sodium hospinin, Water jourfled; Cap removed instructions. To remove cap, squeeze cap at base on opposite sides on the two Jurn impressions with fingers. While squeezing, twist cap countertricokwise to remove. Press down to use. Refer to amove on cap.

To reapply cap, turn cap on pump clockwise until it locks.

"This product is not manufactured or distributed by Bayer HealthCare LLC, the distributor of Afrin" No Drip Severe Congestion Pump Mist.

IMPORTANT: KEEP OUTER CARTON FOR FUTURE REFERENCE ON FULL LABELING

PCVS Health.

Compare to the active ingraciant in Afrin* No Drip Severe Congestion Pump Mist* Sovere

No Drip Nasal (2) Mist

OXYMETAZOLINE HCI 0.05%

Nasal decongestant 12 HOUR PUMP MIST



CVS Quality

LOT



5001014



NO DRIP SEVERE NASAL MIST

oxymetazoline spray liquid

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-315	
Route of Administration	NASAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
OXYMETAZOLINE HYDROCHLORIDE (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZ OLINE HYDROCHLORIDE	50 mg in 100 mL		

Inactive Ingredients		
	Ingredient Name	Strength

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM PHOSPHATE, MONOBASIC (UNII: 3980JIH2SW)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
WATER (UNII: 059QF0KO0R)	
POVIDONE (UNII: FZ989GH94E)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
CAMPHOR (SYNTHETIC) (UNII: 5TJD82A1ET)	
MENTHOL (UNII: L7T10EIP3A)	
EUCALYPTOL (UNII: RV6J6604TK)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842- 315-51	1 in 1 CARTON	08/30/2018	
1		15 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		
2	NDC:69842- 315-30	1 in 1 CARTON	04/30/2022	
2		30 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

Marketing Information			
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/30/2018	

Labeler - CVS Health (062312574)

Registrant - Sheffield Pharmaceuticals LLC (151177797)

Establishment					
Name	Address	ID/FEI	Business Operations		
Sheffield Pharmaceutical LLC		151177797	manufacture(69842-315)		

Revised: 4/2022 CVS Health