

**ASPIRIN REGULAR STRENGTH- aspirin tablet, film coated**  
**FOODHOLD U.S.A., LLC**

-----  
**Care One 44-157**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- asthma (wheezing)
- hives
- facial swelling
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients**

corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?**

**1-800-426-9391**

**Principal Display Panel**

CAREONE®

**NDC 41520-157-87**

*Compare to the  
active ingredient in  
Genuine Bayer® Aspirin<sup>†</sup>*

**Regular Strength  
ASPIRIN**

**325 mg**

Pain Reliever  
Fever Reducer (**NSAID**)

Safe Pain Relief

*Actual  
Size*

**300 TABLETS**

**OUR PHARMACISTS RECOMMEND RX**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

<sup>†</sup>This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.  
50844 REV0122B15717

**DISTRIBUTED BY FOODHOLD U.S.A., LLC  
LANDOVER, MD 20785 • 1-877-846-9949**

**©2022 S&S Brands, LLC • Quality guaranteed or your money back.**

**CAREone**® NDC 41520-157-87  
Compare to the active ingredient in Genuine Bayer® Aspirin®

**Regular Strength**  
**ASPIRIN**

**325 mg**  
Pain Reliever  
Fever Reducer (NSAID)  
Safe Pain Relief

OUR PHARMACISTS RECOMMEND

Actual Size

**300 TABLETS**

**Drug Facts**  
**Active ingredient (in each tablet)** Purpose  
Aspirin 325 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves: ■ minor pain of arthritis ■ muscle pain ■ menstrual pain ■ headache ■ pain and fever of colds ■ toothache

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ asthma (wheezing) ■ hives ■ facial swelling ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed ■ have 3 or more alcoholic drinks every day while using this product

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer  
■ if you have ever had an allergic reaction to this product or any of its ingredients

**Other information:** ■ This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.  
50844 REI0122815717  
DISTRIBUTED BY FOODMOLD U.S.A., LLC  
LANDOVER, MD 20785 • 1-877-846-9849  
©2022 SAS Brands, LLC • Quality guaranteed or your money back.

3 41520-31361 5

**PEEL HERE FOR MORE DRUG FACTS**

**Drug Facts (continued)**

**Ask a doctor before use if** ■ you are taking a diuretic ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have asthma ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ an allergic reaction ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days

**These could be signs of a serious condition.**

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions**  
■ do not take more than directed  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor  
■ children under 12 years: do not use unless directed by a doctor

**Other information** ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

**Inactive ingredients** corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?** 1-800-426-9391

STOP PEELING

CareOne 44-157

## ASPIRIN REGULAR STRENGTH

aspirin tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-157
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg	
Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWIA)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;157;ASPIRIN
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-157-78	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
2	NDC:41520-157-87	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
3	NDC:41520-157-90	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/10/1996	

**Labeler** - FOODHOLD U.S.A., LLC (809183973)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(41520-157)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(41520-157)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(41520-157)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(41520-157)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(41520-157)