

**NYSTATIN- nystatin powder**  
**Northstar Rx LLC**


-----  
**Nystatin Topical Powder, USP**  
**Rx Only**

**For topical use only.**  
**Not for ophthalmic use.**

**DESCRIPTION**

Nystatin is a polyene antifungal antibiotic obtained from *Streptomyces noursei*. The molecular formula for Nystatin is C<sub>47</sub>H<sub>75</sub>NO<sub>17</sub>. The molecular weight of Nystatin is 926.1.

Structural formula:



Nystatin Topical Powder USP is for dermatologic use.

Nystatin Topical Powder USP contains 100,000 USP nystatin units per gram dispersed in talc.

**CLINICAL PHARMACOLOGY**

**Pharmacokinetics**

Nystatin is not absorbed from intact skin or mucous membrane.

**Microbiology**

Nystatin is an antibiotic which is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi, including *Candida albicans*, *C. parapsilosis*, *C. tropicalis*, *C. guilliermondi*, *C. pseudotropicalis*, *C. krusei*, *Torulopsis glabrata*, *Trichophyton rubrum*, *T. mentagrophytes*.

Nystatin acts by binding to sterols in the cell membrane of susceptible species resulting in a change in membrane permeability and the subsequent leakage of intracellular components. On repeated subculturing with increasing levels of nystatin, *Candida albicans* does not develop resistance to nystatin. Generally, resistance to nystatin does not develop during therapy. However, other species of *Candida* (*C. tropicalis*, *C. guilliermondi*, *C. krusei*, and *C. stellatoidea*) become quite resistant on treatment with

nystatin and simultaneously become cross resistant to amphotericin as well. This resistance is lost when the antibiotic is removed. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

## **INDICATIONS AND USAGE**

Nystatin Topical Powder is indicated in the treatment of cutaneous or mucocutaneous mycotic infections caused by *Candida albicans* and other susceptible *Candida* species.

**This preparation is not indicated for systemic, oral, intravaginal or ophthalmic use.**

## **CONTRAINDICATIONS**

Nystatin Topical Powder is contraindicated in patients with a history of hypersensitivity to **any** of its components.

## **PRECAUTIONS**

### **General**

**Nystatin Topical Powder should not be used for the treatment of systemic, oral, intravaginal or ophthalmic infections.**

If irritation or sensitization develops, treatment should be discontinued and appropriate measures taken as indicated. It is recommended that KOH smears, cultures, or other diagnostic methods be used to confirm the diagnosis of cutaneous or mucocutaneous candidiasis and to rule out infection caused by other pathogens.

## **INFORMATION FOR THE PATIENT**

Patients using this medication should receive the following information and instructions:

1. The patient should be instructed to use this medication as directed (including the replacement of missed doses). This medication is not for any disorder other than that for which it is prescribed.
2. Even if symptomatic relief occurs within the first few days of treatment, the patient should be advised not to interrupt or discontinue therapy until the prescribed course of treatment is completed.
3. If symptoms of irritation develop, the patient should be advised to notify the physician promptly.

### **Laboratory Tests**

If there is a lack of therapeutic response, KOH smears, cultures, or other diagnostic methods should be repeated.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

No long-term animal studies have been performed to evaluate the carcinogenic potential of nystatin. No studies have been performed to determine the mutagenicity of nystatin

or its effects on male or female fertility.

### **Pregnancy: Teratogenic Effects**

Category C. Animal reproduction studies have not been conducted with any nystatin topical preparation. It also is not known whether these preparations can cause fetal harm when used by a pregnant woman or can affect reproductive capacity. Nystatin topical preparations should be prescribed for a pregnant woman only if the potential benefit to the mother outweighs the potential risk to the fetus.

### **Nursing Mothers**

It is not known whether nystatin is excreted in human milk. Caution should be exercised when nystatin is prescribed for a nursing woman.

### **Pediatric Use**

Safety and effectiveness have been established in the pediatric population from birth to 16 years. (See **DOSAGE AND ADMINISTRATION**).

## **ADVERSE REACTIONS**

The frequency of adverse events reported in patients using nystatin topical preparations is less than 0.1%. The more common events that were reported include allergic reactions, burning, itching, rash, eczema, and pain on application. (See **PRECAUTIONS: General**.)

## **DOSAGE AND ADMINISTRATION**

Very moist lesions are best treated with the topical dusting powder.

### **Adults and Pediatric Patients (Neonates and Older):**

Apply to candidal lesions two or three times daily until healing is complete. For fungal infection of the feet caused by *Candida* species, the powder should be dusted on the feet, as well as, in all foot wear.

## **HOW SUPPLIED**

Nystatin Topical Powder USP is supplied as 100,000 units nystatin per gram in 15 g, 30 g and 60 g plastic squeeze bottles.

(NDC 72603-491-01)

(NDC 72603-491-02)

(NDC 72603-491-03)

## **STORAGE**

Store at controlled room temperature 15° to 30°C (59° to 86°F); avoid excessive heat (40°C; 104°F).

Manufactured for:  
Northstar Rx LLC  
Memphis, TN 38141

Manufactured by:  
Padagis®  
Minneapolis, MN 55427

Rev. 09-25

## PRINCIPAL DISPLAY PANEL

Rx only

NDC 72603-491-01


Nystatin Topical Powder, USP

100,000 USP Units Per Gram

USUAL DOSAGE: Apply to affected area 2 or 3 times daily.

STORAGE: Store at controlled room temperature 15° to 30°C (59° to 86°); avoid excessive heat (40°C; 104°F).

Net Weight 15 grams



## NYSTATIN

nystatin powder

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:72603-491
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Nystatin</b> (UNII: BDF1O1C72E) (Nystatin - UNII:BDF1O1C72E)	Nystatin	100000 [USP'U] in 1 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>Talc</b> (UNII: 7SEV7J4R1U)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
<b>1</b>	NDC:72603-491-01	15 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/01/2025	
<b>2</b>	NDC:72603-491-02	30 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/01/2025	
<b>3</b>	NDC:72603-491-03	60 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/01/2025	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA064118	09/01/2025	

**Labeler** - Northstar Rx LLC (830546433)

Revised: 9/2025

Northstar Rx LLC