

**EXTRA STRENGTH ACETAMINOHEN ADULT RAPID BURST CHERRY-
acetaminophen liquid
Major Pharmaceuticals**

Drug Facts

Active Ingredients (in each 15 mL)

Acetaminophen 500 mg

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do Not Use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in any 24 hour period
- measure only with dosing cup provided. Do not use any other dosing device
- mL=milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 6 hours while symptoms last
 - do not take more than 10 days unless directed by a doctor
- children under 12 years: do not use

Other Information

- **each 15 mL contains:** sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate
- protect from light

Inactive Ingredients

citric acid, D&C red #33, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

Principal Display Panel

Major **Mapap**

ADULT

For ages 12 years and over

Extra Strength Acetaminophen 500 mg

Rapid Burst Cherry

Pain reliever

fever reducer

*COmpare to Extra strength Tylenol® active ingredient

Alcohol free

FL OZ (mL)

Questions or comments?

Call 1-877-753-3935

Monday-Friday 9AM-5PM EST

Distributed by: MAJOR Pharmaceuticals

17177 N Laurel Park Drive, Suite 233

Livonia, MI 48152 USA

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Extra Strength Tylenol®

Package label



TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Extra Strength Tylenol®.

Drug Facts	
Active ingredient (in each 15 mL)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer

PLD-G2298
LB003856
PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)	
Uses	
<ul style="list-style-type: none"> temporarily relieves minor aches and pains due to <ul style="list-style-type: none"> the common cold backache toothache premenstrual and menstrual cramps temporarily reduces fever headache minor pain of arthritis muscular aches 	
Warnings	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours 	

Drug Facts (continued)	
<ul style="list-style-type: none"> with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin reddening blisters rash If a skin reaction occurs, stop use and seek medical help right away.	
Do not use	
<ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains 	

Drug Facts (continued)	
acetaminophen, ask a doctor or pharmacist.	
Ask a doctor before use if you have liver disease.	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present These could be signs of a serious condition.	

Drug Facts (continued)	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children.	
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
<ul style="list-style-type: none"> do not take more than directed (see Overdose warning) 	

Drug Facts (continued)	
do not take more than 4 doses in any 24-hour period	
<ul style="list-style-type: none"> measure only with dosing cup provided. Do not use any other dosing device. mL = milliliter keep dosing cup with product adults and children 12 years and over <ul style="list-style-type: none"> 30 mL every 6 hours while symptoms last do not take more than 10 days unless directed by a doctor children under 12 years: do not use 	
Other information	
<ul style="list-style-type: none"> each 15 mL contains: sodium 6 mg 	

Drug Facts (continued)	
<ul style="list-style-type: none"> store between 20-25°C (68-77°F). Do not refrigerate. protect from light 	
Inactive ingredients citric acid, D&C red #33, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol	

Major MAPAP

EXTRA STRENGTH ACETAMINOPHEN ADULT RAPID BURST CHERRY acetaminophen liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-5847
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg in 15 mL
Inactive Ingredients			
	Ingredient Name		Strength
	CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)		
	HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)		
	POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ)		
	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
	WATER (UNII: 059QF0KO0R)		
	D&C RED NO. 33 (UNII: 9DBA0SBB0L)		
	FD&C RED NO. 40 (UNII: WZB9127XOA)		
	SACCHARIN SODIUM ANHYDROUS (UNII: I4807BK602)		
	SODIUM BENZOATE (UNII: OJ245FE5EU)		
	SORBITOL (UNII: 506T60A25R)		

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY (Tylenol Type Cherry Rapid Blast Flavor)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-5847-09	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/02/2009	11/28/2026

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/02/2009	11/28/2026

Labeler - Major Pharmaceuticals (191427277)

Revised: 8/2024

Major Pharmaceuticals