

FAMOTIDINE- famotidine tablet, film coated
Ohm Laboratories Inc.

Famotidine

Drug Facts

Active ingredient (in each tablet)

Famotidine, USP 10 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert

Do not use if you are allergic to famotidine or other acid reducers

Do not use

- with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath: sweating: pain spreading to arms, neck or shoulders: or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20° to 25° C (68° to 77° F)
- protect from moisture
- read the directions and warnings before use
- keep the carton. It contains important information.

Inactive ingredients

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

Questions?

call **1-800-406-7984**

Distributed by:

Ohm Laboratories Inc.

1385 Livingston Avenue

North Brunswick, NJ 08902

PRINCIPAL DISPLAY PANEL - 10 mg Tablet Blister Pack Carton

†Compare to the active ingredient of

Original Strength Pepcid AC®

NDC 51660-035-31

ohm®

ORIGINAL STRENGTH

Acid Controller

Famotidine Tablets,

USP 10 mg

Acid Reducer

30 TABLETS

Just One Tablet

Prevents & Relieves Heartburn

Due to Acid Indigestion

Drug Facts

Active Ingredient (in each tablet) Famotidine, USP 10 mg Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Do not use

- if you have trouble or pain swallowing with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- requent chest pain
- requent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Directions

- adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).



ohm®
ORIGINAL STRENGTH
Acid Controller
Famotidine Tablets, USP 10 mg
30 TABLETS Acid Reducer

ohm®
ORIGINAL STRENGTH
Acid Controller
Famotidine Tablets,
USP 10 mg
Acid Reducer

30 TABLETS

Just One Tablet
Prevents & Relieves Heartburn
Due to Acid Indigestion

NDC 51660-035-31

†Compare to the active ingredient of
Original Strength Pepcid AC®



See end panel for batch number and expiration date.

TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.

Ohm® is a registered trademark of Ranbaxy Pharmaceuticals, Inc. All other trademarks are property of their respective owners.

Distributed by:
Ohm Laboratories Inc.
1385 Livingston Avenue
North Brunswick, NJ 08902

R0313


Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Expiration Date: _____
Batch No. _____



JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages. Questions? call 1-800-406-7984



FAMOTIDINE

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51660-035
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (120000 MW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	035
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51660-035-31	1 in 1 CARTON	02/15/2010	
1		30 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090283	02/15/2010	

Labeler - Ohm Laboratories Inc. (184769029)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(51660-035)

Revised: 8/2018

Ohm Laboratories Inc.