RIGHT REMEDIES NIGHTTIME SEVERE COLD AND FLU SOFTGELacetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled Strive Pharmaceuticals Inc.

-----

## RIGHT REMEDIES NIGHTTIME SEVERE COLD & FLU softgel

## **Drug Facts**

## Active ingredients (in each softgel)

Acetaminophen 325 mg Dextromethorphan HBr 10 mg Doxylamine succinate 6.25 mg Phenylephrine HCl 5 mg

## **Purposes**

Pain reliever/fever reducer Cough suppressant Antihistamine Nasal decongestant

#### Uses

- temporarily relieves common cold/flu symptoms:
  - nasal congestion
     sinus congestion
     pressure
  - cough due to minor throat & bronchial irritation
  - cough to help you sleep minor aches & pains
  - headache
     fever
     sore throat
- runny nose & sneezing
   reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 softgels in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have •** liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** • taking sedatives or tranquilizers • taking the blood thinning drug warfarin

## When using this product • do not use more than directed

excitability may occur, especially in children
 marked drowsiness may occur
 alcoholic drinks
 be careful when driving a motor vehicle or operating machinery
 alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** • you get nervous, dizzy, or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

• take only as directed • do not exceed 8 softgels per 24 hrs

adults & children 12	2 softgels with water
yrs & over	every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

#### Other information

• store at controlled room temperature 20–25°C (68–77°F) • protect from light, heat, and moisture

## Inactive ingredients

D&C Yellow No.10, FD&C Blue No.1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide

## Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

Compare to the active ingredients in Vicks<sup>®</sup> NyQuil<sup>™</sup> Severe Cold & Flu LiquiCaps <sup>™</sup>\*

MAX STRENGTH

# TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

\*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks<sup>®</sup> NyQuil™ LiquiCaps™.


Distributed by: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816

PRODUCT OF INDIA

## **Packaging**









■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma
■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers 

taking the blood thinning drug warfarin When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur
■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase drowsiness Stop use and ask a doctor if ■ you get nervous, dizzy, or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headaché that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a
Poison Control Center right away. Quick medical attention is critical for
adults as well as for children even if you do not notice any signs or symptoms. **Directions** ■ take only as directed ■ do not exceed 8 softgels per 24 hrs adults & children 12 yrs & over 2 softgels with water every 4 hrs ask a doctor children 4 to under 12 yrs children under 4 yrs do not use **Other information** ■ store at controlled room temperature 20-25°C (68-77°F) ■ protect from light, heat, and moisture **Inactive ingredients** D&C Yellow No.10, FD&C Blue No.1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide Questions or comments? 1-888-577-8033 Monday - Friday 8am - 4pm EST Distributed by: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816 PRODUCT OF INDIA \_ \_ \_ LB7069283784 REV.00-092025

## RIGHT REMEDIES NIGHTTIME SEVERE COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-837
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

## **Inactive Ingredients**

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6092ICV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	green	Score	no score
Shape	OVAL	Size	18mm
Flavor		Imprint Code	A07
Contains			

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:70692-837- 84	48 in 1 BOTTLE; Type 0: Not a Combination Product	09/23/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/23/2025	

## Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 9/2025 Strive Pharmaceuticals Inc.