ACETAMINOPHEN- acetaminophen tablet, film coated, extended release Medline Industries, LP

643 Acetaminophen

Active ingredient (in each caplet)

Acetaminophen USP, 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - premenstrual and menstrual cramps
 - the common cold
 - headache
 - toothache
- temporarily reduces fever

Warnings

Liver warning:This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed (see overdose warning)

adults	 take 2 caplets every 8 hours with water swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours do not use for more than 10 days unless directed by a doctor
under 18 years of age	ask a doctor

Other information

- store at 20 25° C (68 77° F). Avoid excessive heat 40° C (104° F).
- see end panel for batch number and expiration date
- TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING FROM BOTTLE.

Inactive ingredients

croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, sodium lauryl sulfate, stearic acid, titanium dioxide

Questions?

Call toll free 1-800-MEDLINE (633-5463) Monday-Friday 9AM-5PM CST

Manufacturing Information

Manufactured for:

Medline Industries, LP

Three Lakes Drive, Northfield, IL 60093 USA Made in USA with domestic and foreign materials www.medline.com/www.curad.com 1-800-MEDLINE (633-5463) RFF: OTC33350N

REF: OTC33350N V1 RG220HM

Package Label



Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use with any other drug containing acetaminophen (prescription or non-prescription), if you are not sure whether a drug contains acetaminophen.

Drug Facts (continued)

ask a doctor or pharmacist. III if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have III liver disease.

Ask a doctor or pharmacist before use if you are making the blood thinning drug warfarin. 1

Stop use and ask a doctor If pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days
■ new symptoms occur

m redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued) Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms. Directions on do not take more than directed (see overdose warning) ■ take 2 caplets every 8 hours with water m swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours m do not take for more than 10 days unless directed by a under 18 ask a doctor years of age

Drug Facts (continued)

Other information Store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F)

Inactive ingredients croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, sodium lauryl sulfate, stearic acid, titanium dioxide.

Questions or comments? Call toll free 1-800-MEDLINE (633-5463) Monday-Friday 9AM-5PM CST.

www.medline.com Packaged in the USA with U.S. and foreign components for: Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093 USA. V1 RG220HM 1-800-MEDLINE

REF OTC33350N

Contains No Aspirin

"All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Tylenol® Arthritis Pain.

ACETAMINOPHEN

acetaminophen tablet, film coated, extended release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:53329-643

Route of Administration ORAL

Active Ingredient/Active Moiety

Basis of Strength Strength **Ingredient Name** 650 ma

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D)

ACETAMINOPHEN

Inactive Ingredients			
Ingredient Name	Strength		
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
POVIDONE (UNII: FZ989GH94E)			
STARCH, PREGELATINIZED CORN (UNII: 08232NY3SJ)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
SODIUM LAURYL SULFATE (UNII: 368GB5141J)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

Product Characteristics			
Color	white	Score	no score
Shape	OVAL (Capsule Shaped)	Size	19mm
Flavor		Imprint Code	cor116
Contains			

l	Packaging				
#		# Item Code Package Description		Marketing Start Date	Marketing End Date
		NDC:53329-643- 29	50 in 1 BOTTLE; Type 0: Not a Combination Product	07/10/2018	12/31/2025

Marketing I	nformation		
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076200	04/30/2002	12/31/2025

Labeler - Medline Industries, LP (025460908)

Registrant - Medline Industries, LP (025460908)

Establishment					
Name	Address	ID/FEI	Business Operations		
Ohm Laboratories Inc.		184769029	manufacture(53329-643)		

Revised: 1/2025 Medline Industries, LP