REGULAR STRENGTH ANTACID- aluminum hydroxide, magnesium hydroxide, dimethicone suspension Chain Drug Marketing Association

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

qc ant org

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 200 mg (equivalent to dried gel, USP)

Magnesium hydroxide 200 mg

Simethicone 20mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- do not take more than 24 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks

• children under 12 years: ask a doctor

Other information

- each 5 mL teaspoonful contains: magnesium 85 mg, sodium 3 mg
- store at room temperature
- protect from freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or Comments?

1-800-540-3765

package Label

NDC 63868-712-57



Compare to the Active Ingredients in Mylanta Regular Strength

Regular Strength Antacid

Antigas, Fast Acting

Alumina, Magnesia, and Simethicone Oral Suspension USP

Fast Soothing Relief of:

Acid Indigestion Heartburn Sour Stomach Pressure & Bloating

Alcohol: 0.15%

Original Flavor

12 FL OZ (355 mL)

REV GC-0818 231-05112-0

Drug Facts

TAMPER-EVIDENT: Do not use if breakaway band on bottle cap is missing or broken.

Uses relieves • heartburn • sour stomach • acid indigestion
 • the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

· kidney disease · a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last for more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions • shake well before use

- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls between meals, at bedtime or as directed by a doctor
- do not take more than 24 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor

Other information • each 5 mL teaspoonful contains:

magnesium 85 mg, sodium 3 mg • keep tightly closed

protect from freezing

store at room temperature

Inactive ingredients benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Ouestions or comments? 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark MYLANTA®.



REV GC-0818 931-05112-0

Distributed by C.D.M.A., Inc.® 43157 W. Nine Mile Novi, MI 48376-0995 www.qualitychoice.com Questions: 248-449-9300



REGULAR STRENGTH ANTACID

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:63868-712 Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
AL LIMITED AND COURSE (LIMIT CODOTTOLINO) (AL LIMITED DOVIDE LIMIT CODOTTOLINO)	ALUMINUM	200 mg		

ALUMINUM III DRU AIDE (UMI. 5Q)	DO 1210100) (MEGIMINOIMITITUMOMIDE - OMILIOQUO 1210110	' HYDRO XIDE	in 5 mL
MAGNESIUM HYDRO XIDE (UNII: NI HYDRO XIDE ION - UNII:9 159 UV38 1P	BZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838,)	MAGNESIUM HYDRO XIDE	200 mg in 5 mL
DIMETHICO NE (UNII: 92RU3N3Y10)) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 5 mL

Inactive Ingredients				
Ingredient Name	Strength			
BENZYL ALCOHOL (UNII: LKG8494WBH)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
HYDROXYETHYL CELLULOSE (4000 MPA.S AT 1%) (UNII: ZYD53NBL45)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)				
SORBITOL (UNII: 506T60A25R)				

Product Characteristics				
Color		Score		
Shape		Size		
Flavor	LEMON (citrus mint)	Imprint Code		
Contains				

Packaging					
l	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
l	1 I	NDC:63868-712-57	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2018	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part331	03/01/2014		

Labeler - Chain Drug Marketing Association (011920774)

Registrant - GCP Laboratories (965480861)

Establishment			
Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(63868-712)

Revised: 12/2018 Chain Drug Marketing Association