

BASIC CARE DAYTIME SEVERE COLD AND FLU NIGHTTIME SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, doxylamine succinate
Amazon.com Services LLC

Amazon Daytime Severe Cold & Flu Nighttime Severe Cold & Flu Drug Facts

Active ingredients (in each caplet) Daytime

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- fever
- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- headache
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- thyroid disease
- diabetes
- liver disease
- heart disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see overdose warning
- do not exceed 8 caplets per 24 hrs

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each caplet contains:** sodium 3 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, flavor, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Active ingredients (in each caplet) Nighttime

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see overdose warning
- do not exceed 8 caplets per 24 hrs

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at 20-25° C (68-77° F)

Inactive ingredients

crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Daytime Relief

Compare to Vicks® DayQuil® Severe+ VapoCOOL™

active ingredients

Non-Drowsy

Daytime Severe Cold & Flu

Acetaminophen

Phenylephrine HCl

Dextromethorphan HBr

Guaifenesin

actual size

Pain Reliever/Fever Reducer

Nasal Decongestant

Cough Suppressant

Expectorant

Max Strength

16 CAPLETS

Vapor Ice™

Nighttime Relief

Compare to Vicks® NyQuil® Severe+ VapoCOOL™

active ingredients

Nighttime Severe Cold & Flu

Acetaminophen

Phenylephrine HCl

Doxylamine Succinate

Dextromethorphan HBr

actual size

Pain Reliever/Fever Reducer

Nasal Decongestant

Antihistamine

Cough Suppressant

Max Strength

8 CAPLETS

Vapor Ice™

amazon

basic+care

Daytime
Relief

Compare to
Vicks® DayQuil®
Severe+
VapoCOOL™
active ingredients*

Non-Drowsy

Daytime Severe Cold & Flu

Acetaminophen
Phenylephrine HCl
Dextromethorphan HBr
Guaifenesin

Pain Reliever/Fever Reducer
Nasal Decongestant
Cough Suppressant
Expectorant

Max Strength

actual
size



Vapor Ice™

16 CAPLETS

Nighttime
Relief

Compare to
Vicks® NyQuil®
Severe+
VapoCOOL™
active ingredients*

NDC 72288-002-90

Nighttime Severe Cold & Flu

Acetaminophen
Phenylephrine HCl
Doxylamine Succinate
Dextromethorphan HBr

Pain Reliever/Fever Reducer
Nasal Decongestant
Antihistamine
Cough Suppressant

Max Strength

actual
size



Vapor Ice™


8 CAPLETS

Drug Facts	Nighttime SEVERE+ Vapor Ice™ Cold & Flu	Drug Facts	Daytime SEVERE+ Vapor Ice™ Cold & Flu
Active ingredients (in each caplet) Acetaminophen 325 mg Dextromethorphan HBr 10 mg Doxylamine succinate 6.25 mg Phenylephrine HCl 5 mg	Purpose Pain reliever/fever reducer Cough suppressant Antihistamine Nasal decongestant	Active ingredients (in each caplet) Acetaminophen 325 mg Dextromethorphan HBr 10 mg Guaifenesin 200 mg Phenylephrine HCl 5 mg	Purpose Pain reliever/fever reducer Cough suppressant Expectorant Nasal decongestant
Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ cough that upsets your sleep ■ minor aches and pains ■ headache ■ fever ■ sore throat ■ runny nose and sneezing ■ reduces swelling of nasal passages ■ temporarily restores free breathing through the nose ■ promotes nasal and/or sinus drainage		Uses temporarily relieves common cold/flu symptoms: ■ fever ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ headache ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores free breathing through the nose ■ minor aches and pains ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passages ■ helps loosen mucus and make acute coughs more productive ■ promotes nasal and/or sinus drainage	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs		Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs	

921.90 BN C1

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

*These products are not manufactured or distributed by Procter & Gamble, distributor of Wicks® Daytime® Severe+ Vapor Ice™ and Wicks® Nighttime® Severe+ Vapor Ice™. GLUTEN FREE



3 700301 14526 3

Drug Facts (continued)

containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ high blood pressure ■ thyroid disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as asthma or chronic bronchitis ■ persistent or chronic cough such as occurs with asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ thyroid disease ■ diabetes ■ liver disease ■ heart disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts (continued)
Inactive ingredients: croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, flavor, malic acid, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? 1-800-719-9280

DISTRIBUTED BY:
AMA-ZOLA.COM SERVICES LLC
410 TERRY AVENUE N., SEATTLE, WA 98109

Drug Facts (continued)

Directions ■ take only as directed - see overdose warning
■ do not exceed 8 caplets per 24 hrs

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, flavin magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? 1-800-719-9280

Drug Facts (continued)

Directions ■ take only as directed - see overdose warning
■ do not exceed 8 caplets per 24 hrs

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each caplet contains sodium 3 mg ■ store at 20-25°C (68-77°F)

BASIC CARE DAYTIME SEVERE COLD AND FLU NIGHTTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, doxylamine succinate kit

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)		NDC:72288-002
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-002-90	1 in 1 CARTON; Type 0: Not a Combination Product	02/04/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	8 BLISTER PACK	16
Part 2	4 BLISTER PACK	8

Part 1 of 2

BASIC CARE DAYTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride tablet, film coated

Product Information

Item Code (Source)	NDC:72288-357
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (120 .MU.M) (UNII: 68401960MK)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	L35C
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-357-00	2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Part 2 of 2

BASIC CARE NIGHTTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride tablet, film coated

Product Information

Item Code (Source)	NDC:72288-721
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CROSPVIDONE (120 .MU.M) (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	GREEN	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	L72V
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-721-00	2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/04/2021	

Labeler - Amazon.com Services LLC (128990418)