

ANTISEPTIC SKIN CLEANSER- chlorhexidine gluconate solution
Xttrium Laboratories, Inc.

Dyna-Hex 4: 4oz FM, 8oz, 16oz, 32oz, 1 Gallon
COMP 18ozFM

Active Ingredient

Chlorhexidine gluconate

Drug Facts

Purpose

Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **skin wound and general skin cleansing**

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area
- as a preoperative skin preparation of the head or face (only in 4-oz)

When using this product

- keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Direction

- use with care in premature infants and infants under 2 months of age. These

products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 ml of product and a wet brush paying close attention to the nails, cuticles and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 ml of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 ml of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Package label and principal display panel

NDC 0116-1061-08

DYNA-HEX 4 ®

(Chlorhexidine Gluconate 4% Solution)

Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed by: Xttrium Laboratories, Inc., Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 8 fl oz (237 mL)

4DY08BTLLBLF

NDC 0116-1061-08	
DYNA-HEX 4[®] Chlorhexidine Gluconate 4% Solution <i>Antiseptic</i>	
Contains: 4% Chlorhexidine Gluconate Distributed By: Xttrium Laboratories, Inc. Mount Prospect, IL 60056 Net Contents: 8 fl oz (237 mL)	
Drug Facts	Drug Facts (continued)
Active ingredient Chlorhexidine gluconate 4% solution Antiseptic	Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: ■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash If an allergic reaction occurs, stop use and seek medical help right away.
Uses ■ surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care ■ healthcare personnel handwash: helps reduce bacteria that potentially can cause disease ■ skin wound and general skin cleansing	Do not use ■ if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product ■ in contact with meninges ■ in the genital area
Warnings For external use only	When using this product ■ keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums. ■ if solution should contact these areas, rinse out promptly and thoroughly with water ■ wounds which involve more than the superficial layers of the skin should not be routinely treated
	Drug Facts (continued) ■ repeated general skin cleansing of large body areas should not be done except when advised by a health care provider Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
	Directions ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Surgical hand scrub: ■ wet hands and forearms with water ■ scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces ■ a separate nail cleaner may be used ■ rinse thoroughly ■ wash for an additional 3 minutes with 5 mL of product and rinse under running water ■ dry thoroughly
	Drug Facts (continued) Healthcare personnel handwash: ■ wet hands with water ■ dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds ■ rinse and dry thoroughly Skin wound and general skin cleaning: ■ thoroughly rinse the area to be cleaned with water ■ apply the minimum amount of product necessary to cover the skin or wound area and wash gently ■ rinse again thoroughly
	Other information ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)
	Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol
	Questions or comments? call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundry/Cleaning Instructions:
Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

3 01161 06108 5
NDC 01161-1061-08
4DY08BTLLBLF

NDC 0116-1061-16

DYNA-HEX 4[®]

(Chlorhexidine Gluconate 4% Solution)

Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed by: Xttrium Laboratories, Inc., Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 16 fl oz (473 mL)

4DY16BTLLBLF

NDC 0116-1061-16

DYNA-HEX 4[®]

Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate
Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056
Net Contents: 16 fl oz (473 mL)

Drug Facts

Active ingredient Chlorhexidine gluconate 4% solution
Purpose Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **skin wound and general skin cleansing**

Warnings

For external use only

Drug Facts (continued)

Allergy alert:
This product may cause a severe allergic reaction. Symptoms may include:
■ wheezing/difficulty breathing
■ shock ■ facial swelling
■ hives ■ rash
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meningitis
- in the genital area

When using this product

- keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated

Drug Facts (continued)

- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Drug Facts (continued)

Healthcare personnel handwash:

- wet hands with water
 - dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
 - rinse and dry thoroughly
- Skin wound and general skin cleaning:**
- thoroughly rinse the area to be cleaned with water
 - apply the minimum amount of product necessary to cover the skin or wound area and wash gently
 - rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundry/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.



NDC 0116-1061-32

DYNA-HEX 4[®]

(Chlorhexidine Gluconate 4% Solution)

Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed by: Xttrium Laboratories, Inc., Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 32 fl oz (946 mL)

4DY32BTLBLB

NDC 0116-1061-32

DYNA-HEX 4[®]

Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate
Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056



FOR EXTERNAL USE ONLY

Net Contents: 32 fl oz (1 qt) (946 mL)

4DY32BTLBLB

Drug Facts

Active ingredient Chlorhexidine gluconate 4% solution
Purpose Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **skin wound and general skin cleansing**

Warnings

For external use only

Allergy alert:
This product may cause a severe allergic reaction. Symptoms may include:
■ wheezing/difficulty breathing
■ shock ■ facial swelling
■ hives ■ rash
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient are allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meningitis
- in the genital area

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
 - dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
 - rinse and dry thoroughly
- Skin wound and general skin cleaning:**
- thoroughly rinse the area to be cleaned with water
 - apply the minimum amount of product necessary to cover the skin or wound area and wash gently
 - rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol.

Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundry/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

NDC 0116-1061-01

DYNA-HEX 4[®]

(Chlorhexidine Gluconate 4% Solution)

Antiseptic

Contains: 4% Chlorhexidine Gluconate

Distributed by: Xttrium Laboratories, Inc., Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 128 fl oz (1 gal)

4DYN1GBTLLBLC

<p>Drug Facts (continued)</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <p>Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</p> <p>Surgical hand scrub:</p> <ul style="list-style-type: none">Wet hands and forearms with water.Scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces.A separate nail cleaner may be used.Rinse thoroughly.Wash for an additional 3 minutes with 5 mL of product and rinse under running water.Rinse thoroughly. <p>Healthcare personnel handwash:</p> <ul style="list-style-type: none">Wet hands with water.Dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds.Rinse and dry thoroughly. <p>Skin wound and general skin cleansing:</p> <ul style="list-style-type: none">Thoroughly rinse the area to be cleaned with water.Apply the minimum amount of product necessary to cover the skin or wound area and wash gently.Rinse again thoroughly. <p>Other information</p> <ul style="list-style-type: none">Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). <p>Inactive ingredients: cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, isosorbimide solids, PEG-75 lauryl, purified water, triethyl alcohol.</p> <p>Questions or comments? call 1-800-587-3721 Monday through Friday 8:00 AM to 4:00 PM CST</p> <p><small>Laundry/Cleaning Instructions: Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.</small></p>	<p>NDC 0116-1061-01</p> <p>DYNA-HEX 4[®] Chlorhexidine Gluconate 4% Solution <i>Antiseptic</i></p> <p>Contains: 4% Chlorhexidine Gluconate Distributed By: Xttrium Laboratories, Inc. Mount Prospect, IL 60056</p>  <p>0 11610 61010 7</p> <p>CHLORHEXIDINE GLUCONATE</p> <p>Net Contents: 128 fl oz (1 gal) (3.785 L) 4DYN1GBTLLBLC</p>	<p>Drug Facts</p> <p>Active ingredient Chlorhexidine gluconate 4% solution</p> <p>Purpose Antiseptic</p> <p>Uses</p> <ul style="list-style-type: none">Surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care.Healthcare personnel handwash: helps reduce bacteria that potentially can cause disease.Skin wound and general skin cleansing. <p>Warnings</p> <p>For external use only</p> <p>Allergy alert: This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none">itching/difficulty breathingswelling of the face, lips, tongue, or throathivesfacial swellingrash <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use</p> <ul style="list-style-type: none">if you or the patient are allergic to chlorhexidine gluconate or any other ingredient in this productin contact with mucous membranesin the genital area <p>When using this product</p> <ul style="list-style-type: none">keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause blindness when inhaled in the middle ear through perforated eardrums.if children should contact their ears, rinse out promptly and thoroughly with water.wounds which involve more than the superficial layers of the skin should not be routinely treated.repeated genital skin cleansing of large body areas should not be done except when advised by a health care provider. <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p>
---	--	--

NDC 0116-1061-40

DYNA-HEX 4[®]

Chlorhexidine Gluconate 4% Solution

Antiseptic

Contains:

4% Chlorhexidine Gluconate With foam pump

Distributed By:

Xttrium Laboratories, Inc.

Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 4 fl oz (118mL)

4DY04FM1061LBLB

NDC 0116-1061-40

DYNA-HEX 4®

Chlorhexidine Gluconate 4%
Solution Antiseptic

Contains: 4% Chlorhexidine Gluconate
With foam pump
Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056
Net Contents: 4 fl oz (118 mL)

Drug Facts

Active ingredient Purpose
Chlorhexidine gluconate 4% solution Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **skin wound and general skin cleansing**

Warnings

For external use only

Drug Facts (continued)

Allergy alert:
This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- hives
- facial swelling
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meningies
- in the genital area

When using this product

- keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be

Drug Facts (continued)

done except when advised by a healthcare provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- Surgical hand scrub:**
- wet hands and forearms with water
 - scrub for 3 minutes with 4 full pumps (6 grams) of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces
 - a separate nail cleaner may be used
 - rinse thoroughly
 - wash for an additional 3 minutes with 4 full pumps (6 grams) of product and rinse under running water
 - dry thoroughly

Drug Facts (continued)

Healthcare personnel handwash:

- wet hands with water
- dispense 4 full pumps (6 grams) of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Skin wound and general skin cleansing:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundry/Cleaning Instructions:
Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.



NDC 0116-1061-18

DYNA-HEX 4®

Chlorhexidine Gluconate 4% Solution

Antiseptic

Contains:

4% Chlorhexidine Gluconate

Distributed By:

Xttrium Laboratories, Inc.

Mount Prospect, IL 60056

FOR EXTERNAL USE ONLY

Net Contents: 18 fl oz (532mL)

4DY18FM1061LBLE

NDC 0116-1061-18

DYNA-HEX 4®

Chlorhexidine Gluconate 4% Solution
Antiseptic

Contains: 4% Chlorhexidine Gluconate
With foam pump
Distributed By: Xttrium Laboratories, Inc.
Mount Prospect, IL 60056
Net Contents: 18 fl oz (532 mL)



Drug Facts

Active ingredient Purpose
Chlorhexidine gluconate 4% solution Antiseptic

Uses

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease
- **skin wound and general skin cleansing**

Warnings

For external use only

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

Drug Facts (continued)

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in the product
- in contact with the meningies
- in the genital area

When using this product

- keep out of eyes, ears and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large

Drug Facts (continued)

body areas should not be done except when advised by a healthcare provider

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- Surgical hand scrub:**
- wet hands and forearms with water
 - scrub for 3 minutes with 4 full pumps (about 5 mL) of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces

Drug Facts (continued)

- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 4 full pumps (about 5 mL) of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense 4 full pumps (about 5 mL) of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Skin wound and general skin cleansing:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Drug Facts (continued)

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments?

call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

Laundry/Cleaning Instructions:
Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents.

ANTISEPTIC SKIN CLEANSER

chlorhexidine gluconate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0116-1061
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	4 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
HYDROXYETHYL CELLULOSE (2000 CPS AT 1%) (UNII: S38J6RZN16)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
LAURAMINE OXIDE (UNII: 4F6FC4MI8W)	
WATER (UNII: 059QF0KO0R)	
TRIDECYL ALCOHOL (UNII: 8I9428H868)	
GLUCONOLACTONE (UNII: WQ29KQ9POT)	
PEG-75 LANOLIN (UNII: 09179OX7TB)	
COCO DIETHANOLAMIDE (UNII: 92005F972D)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0116-1061-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	
2	NDC:0116-1061-16	437 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	
3	NDC:0116-1061-32	946 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	
4	NDC:0116-1061-01	15142 mL in 1 JUG; Type 0: Not a Combination Product	08/31/2011	
5	NDC:0116-1061-40	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2019	
6	NDC:0116-1061-18	532 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/31/2011	08/31/2026

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019125	08/31/2011	

Labeler - Xttrium Laboratories, Inc. (007470579)

Revised: 12/2025

Xttrium Laboratories, Inc.