

ASPI-COR- aspirin tablet, delayed release
L.N.K. International, Inc.

Aspi-Cor 44-675

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- new symptoms occur
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- use by expiration date on package

Inactive ingredients

black iron oxide, colloidal silicon dioxide, corn starch, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, polydextrose, polyethylene glycol, propylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

844-232-8960

Principal display panel

Aspi-Cor 81 mg[®]

Adult Low Dose Aspirin 81 mg
Pain Reliever **(NSAID)**

ENTERIC

**SAFETY
COATED**

120 Tablets

Actual Size

Distributed by:
HEALTH PRODUCTS CORPORATION
Yonkers, N.Y. 10703 NDC 50844-675-32
Product #6ASPI81 REV0422F67532

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

Aspi-Cor 81 mg[®]

Adult Low Dose Aspirin 81 mg
Pain Reliever (NSAID)

ENTERIC

SAFETY
COATED

120 Tablets Actual Size



Distributed by:
HEALTH PRODUCTS CORPORATION
Yonkers, N.Y. 10703 NO: 508-44-675-32
Product #ASPI81 REV/04/22/67532



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purpose

Aspirin 81 mg (NSAID) * Pain reliever
*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

GTIN:00039686014926

EXP.:xxxxxxx

LOT:XXXXXXX

FPO
D2

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding: ■ vomit blood

■ have bloody or black stools ■ feel faint

■ have stomach pain that does not get better

■ ringing in the ears or a loss of hearing occurs

■ new symptoms occur

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

These could be signs of a serious condition.

STOP PEELING

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

black iron oxide, colloidal silicon dioxide, corn starch, croscarmellose sodium, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, polydextrose, polyethylene glycol, propylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

844-232-8960

Aspi-Cor 44-675

ASPI-COR

aspirin tablet, delayed release

Product Information						
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:50844-675		
Route of Administration		ORAL				
Active Ingredient/Active Moiety						
Ingredient Name			Basis of Strength	Strength		
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)			ASPIRIN	81 mg		
Inactive Ingredients						
Ingredient Name				Strength		
FERROSOFERRIC OXIDE (UNII: XM0M87F357)						
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)						
STARCH, CORN (UNII: O8232NY3SJ)						
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)						
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)						
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)						
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)						
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)						
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)						
POLYDEXTROSE (UNII: VH2XOU12IE)						
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)						
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)						
SHELLAC (UNII: 46N107B71O)						
DIMETHICONE (UNII: 92RU3N3Y1O)						
WATER (UNII: 059QF0KO0R)						
SODIUM BICARBONATE (UNII: 8MDF5V39QO)						
SODIUM LAURYL SULFATE (UNII: 368GB5141J)						
TALC (UNII: 7SEV7J4R1U)						
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)						
TRIACETIN (UNII: XHX3C3X673)						
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)						
Product Characteristics						
Color	yellow	Score	no score			
Shape	ROUND	Size	7mm			
Flavor		Imprint Code	ASPI;COR			
Contains						
Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:50844-675-03	10 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/03/2015			
2	NDC:50844-675-32	120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/03/2015			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/03/2015	

Labeler - L.N.K. International, Inc. (038154464)

Revised: 8/2024

L.N.K. International, Inc.