

**COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen, dextromethorphan hbr, phenylephrine hcl, doxylamine succinate
WinCo Foods, LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients in Daytime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Active ingredients in Nighttime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose for Daytime

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Purpose for Nighttime

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - cough due to minor throat and bronchial irritation
 - nasal congestion (Daytime only)

- runny nose and sneezing (Nighttime only)

Warnings

Liver warning: These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease (Daytime-Nighttime)
- heart disease (Daytime)
- diabetes (Daytime)
- thyroid disease (Daytime)
- high blood pressure (Daytime)
- glaucoma (nighttime)
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, emphysema (Daytime)
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema (Nighttime)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Nighttime only)

When using this product,

Daytime: do not use more than directed.

Nighttime:

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur (Daytime only)
- pain, cough, and nasal congestion gets worse or lasts more than 7 days (Daytime)
- pain or cough gets worse or lasts more than 3 days (Nighttime)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve

Daytime: adults and children 12 years and over: take 2 softgels with water every 4 hours

Nighttime: adults and children 12 years and over: take 2 softgels with water every 6 hours

- children under 12 years: do not use

Other information

- store at room temperature 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients for Daytime

butylated hydroxyanisole, butylated hydroxytoluene, carminic acid*, D&C yellow #10*, edible white ink, FD&C red #40*, FD&C yellow #6*, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sodium metabisulfite*, sorbitan, sorbitol

*contains one or more of these ingredients

Inactive ingredients in Nighttime

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan*, sorbitol

*may contain this ingredient

Questions or comments?

Or to report an adverse event call **800-824-1706** Monday-Friday 9am-4pm MST

Principal Display Panel

**Compare to the Active Ingredients in VICKS® DAYQUIL® & NyQUIL® LiquiCaps®

Multi-Symptom Cold & Flu Relief

NON-DROWSY

DAYTIME

ACETAMINOPHEN / PAIN RELIEVER - FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

PHENYLEPHRINE HCl / NASAL DECONGESTANT

Relieves: Aches, Fever, Sore Throat, Cough & Nasal Congestion

Nighttime

ACETAMINOPHEN / PAIN RELIEVER - FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE / ANTIHISTAMINE

Relieves: Aches, Fever, Sore Throat, Cough, Sneezing & Runny Nose.

Softgels

**Liquid Filled Capsules

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS

TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

†This product is not manufactured or distributed by THE PROCTER & GAMBLE COMPANY, VICKS®, DAYQUIL®, NYQUIL® AND LiquiCap® are registered trademarks of THE PROCTER & GAMBLE COMPANY.

DISTRIBUTED BY:

WINCO FOODS, LLC

BOISE, ID 83704

www.wincofoods.com

Product Label

<p>Other Information</p> <ul style="list-style-type: none"> ■ store between 15-30°C (59-86°F) ■ avoid excessive heat ■ children under 12 years: do not use ■ adults and children 12 years and over: take 2 softgels with water every 6 hours. ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve 	<p>Other Information</p> <ul style="list-style-type: none"> ■ store between 15-30°C (59-86°F) ■ avoid excessive heat ■ children under 12 years: do not use ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve
<p>Directions</p> <p>Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Signs of a serious condition:</p> <ul style="list-style-type: none"> ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleepiness occur 	<p>Directions</p> <p>Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Signs of a serious condition:</p> <ul style="list-style-type: none"> ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleepiness occur
<p>Stop use and ask a doctor if</p> <p>When using this product, do not use more than directed.</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <ul style="list-style-type: none"> ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ liver disease ■ heart disease 	<p>Stop use and ask a doctor if</p> <p>When using this product, do not use more than directed.</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <ul style="list-style-type: none"> ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ thyroid disease ■ high blood pressure ■ liver disease ■ heart disease

<p>Drug Facts (continued)</p> <p>Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product, do not use more than directed.</p>	<p>Drug Facts (continued)</p> <p>Ask a doctor before use if you have ■ liver disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product, do not use more than directed.</p>
<p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	<p>Warnings</p> <p>Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>
<p>Uses ■ temporarily relieves common cold and flu symptoms ■ cough due to minor throat and bronchial irritation ■ nasal congestion ■ minor aches and pains ■ fever ■ sore throat</p> <p>Active ingredients (in each softgel)</p> <p>Acetaminophen 325 mg ■ Pain reliever/fever reducer Dextromethorphan HBr 10 mg ■ Cough suppressant Phenylephrine HCl 5 mg ■ Nasal decongestant Antihistamine</p>	<p>Uses ■ temporarily relieves common cold and flu symptoms ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing</p> <p>Active ingredients (in each softgel)</p> <p>Acetaminophen 325 mg ■ Pain reliever/fever reducer Dextromethorphan HBr 15 mg ■ Cough suppressant Doxylamine succinate 6.25 mg ■ Antihistamine</p>

WinCo FOODS

MULTI-SYMPTOM Cold & Flu Relief

Total **48** Softgels**

**Liquid Filled Capsules

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing

Compare to the active ingredients in VICKS® DAYQUIL® & NYQUIL® Liquid Capsules*

NON-DROWSY

Daytime

**ACETAMINOPHEN/
PAIN RELIEVER - FEVER REDUCER**

**DEXTROMETHORPHAN HBr/
COUGH SUPPRESSANT**

**PHENYLEPHRINE HCl/
NASAL DECONGESTANT**

Relieves: Aches, Fever, Sore Throat, Cough & Nasal Congestion



32 Softgels**

**Liquid Filled Capsules

Nighttime

**ACETAMINOPHEN/
PAIN RELIEVER - FEVER REDUCER**

**DEXTROMETHORPHAN HBr/
COUGH SUPPRESSANT**

**DOXYLAMINE SUCCINATE/
ANTIHISTAMINE**

Relieves: Aches, Fever, Sore Throat, Cough, Sneezing & Runny Nose



16 Softgels**

**Liquid Filled Capsules

WinCo FOODS

MULTI-SYMPTOM Cold & Flu Relief

Nighttime

16 Softgels**

**Liquid Filled Capsules

Compare to the active ingredients in VICKS® DAYQUIL® & NYQUIL® Liquid Capsules*

NON-DROWSY

Daytime

**ACETAMINOPHEN/
PAIN RELIEVER - FEVER REDUCER**

**DEXTROMETHORPHAN HBr/
COUGH SUPPRESSANT**

**PHENYLEPHRINE HCl/
NASAL DECONGESTANT**

Relieves: Aches, Fever, Sore Throat, Cough & Nasal Congestion



32 Softgels**

**Liquid Filled Capsules

Nighttime

**ACETAMINOPHEN/
PAIN RELIEVER - FEVER REDUCER**

**DEXTROMETHORPHAN HBr/
COUGH SUPPRESSANT**

**DOXYLAMINE SUCCINATE/
ANTIHISTAMINE**

Relieves: Aches, Fever, Sore Throat, Cough, Sneezing & Runny Nose



16 Softgels**

**Liquid Filled Capsules



Comments & Questions? 800-824-1706 www.wincofoods.com PLD-842R FC03002 WFL411

Lot / Exp.: DT Product of / NT Product of:

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	P30;94A;215;P120
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		16 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/30/2016	05/31/2024

Part 2 of 2

COLD AND FLU RELIEF MULTI SYMPTOM NON DROWSY DAYTIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
CARMINIC ACID (UNII: CID8Z8N95N)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	P19;95A;512;P119
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		32 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC monograph final	part341	09/30/2016	05/31/2024

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/30/2016	12/31/2024

Labeler - WinCo Foods, LLC (056098817)

Revised: 10/2023

WinCo Foods, LLC