UPSET STOMACH RELIEF TO GO- bismuth subsalicylate powder Breakthrough Products Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Upset Stomach Relief to Go™

Drug Facts

Active ingredient (per powder)

Bismuth subsalicylate 525mg

Purpose

Upset stomach reliever/antidiarrheal

Uses

Relieves:

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Contains salicylate.

Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for:

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts for more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years and over:

- see instructions in red box for opening packet
- place 1 powder on the tongue and swallow with or without water every ½ to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

children under 12 years: ask a doctor

Other information

each dose contains salicylate 199mg

Inactive ingredients

citric acid, flavor, glucose, silica, sucralose

Questions?

1-888-998-7436 (Mon-Fri 9AM-5PM MDT)

Do not use if pouch is torn or open

distributed by URGENT Rx®
Breakthrough Products, Inc.
Denver, CO 80202

PRINCIPAL DISPLAY PANEL - 1 Powder Packet

New!

NO

WATER

REQUIRED

POUR DIRECTLY IN MOUTH

 $URGENT Rx^{\mathbb{R}}$

FAST

 $POWDERS_{\text{TM}}$

 $\ \square$ SEE OPENING INSTRUCTIONS ON BACK PANEL $\ \square$

UPSET

STOMACH

 $URGENT Rx^{\mathbb{R}}$

RELIEF

 $TO\text{-}GO_{\mathsf{TM}}$

BISMUTH SUBSALICYLATE/

UPSET STOMACH RELIEVER & ANTIDIARRHEAL

cherry

distributed by UrgentRx 1 POWDER PACK







US Pat. No 8,376,140 CCUS0714-A

 \bigcirc

Do not use if pouch is torn or open

Upset Stomach Relief to Go"

URGENT R®

Breakthrough Products, Inc.

Fold top and store in purse or wallet

Drug Facts

) WHILE FOLDED, TEAR AT NOTCH (or use scissors)

FOLD ON LINE TO EXPOSE NOTCH

Active Ingredient (per powder)

Bismuth subsaicylate 525mg......Upset atomach reliever/antidiamheal

Lees Relieves: • traveler's diarrhea • diamhea • upset atomach due
to overindulgence in food and drink, including: • heartburn
• indigeration • nausea • gas • belching • fullness Warnings

Reye's syndrome: Children and teenagers who have or are
recovering from chickenpox or flu-like symptoms should not use this
product. When using this product, if changes in behavior with nausea
and vomiting occur, consult a doctor because these symptoms could
be an early sign of Reye's syndrome, a rare but serious illness.

Altergy alert: Contains salicylate. Do not take it you are • allergic

Altergy alert: Contains salicylate. Do not take it you are • allergic

to salicylates (including aspirin) • taking other salicylate products Do

not use if you have • an ulcer • a bleeding problem • bloody or black stool Ask a doctor or doctor before use if you have • fever • mucus in the stool Ask a doctor or pharmacist before use if you are taking any drug for: • anticoagulation (thinning the blood) • diabetes • gout • arthritis When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur. Stop use and ask a doctor occurs • diames get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diames lasts for more than 2 days • ringing in the ears of overdose, get health protessional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Directions adults and medical help or contact a Poison Control Center right away. Directions adults and on the tongue and awallow with or without water every ½ to 1 hour as needed • place 1 and the tongue and swallow with or without water every ½ to 1 hour as needed • place 1 and a stop of clear fluids to help prevent dehydration caused by diarrhea children 2 days • under 12 years and over: • see instructions in red box for opening packet • place 1 and one of clear fluids to help prevent dehydration caused by diarrhea children and one exceed 8 doces in 24 hours • use until diarrhea stops but not more than 2 days • under 12 years: ask a doctor Other Information each day diarrhea contains salicylate under 12 years: ask a doctor Other information each dose contains salicylate under 12 years; ask a doctor Other information, glucose, silica, sucralose under on the contains and one ingented of the sucralose of or or described ingented in a doctor Other and one of one or described in a doctor of the and one of or described in a doctor of the and of the and of or described in a doctor of the and of the

(TOM M92-M46 h9-noM) 8647-866-888-1 (snoissau)



■SEE OPENING INSTRUCTIONS ON BACK PANEL ▶

JPSET STOMACH



BISMUTH SUBSALICYLATE/

cherry

distributed by UrgentRx

1 POWDER PACK

UPSET STOMACH RELIEF TO GO

bismuth subsalicylate powder

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51596-004
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Bismuth subsalicylate (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD118YE)	Bismuth subsalicylate	525 mg	

Inactive Ingredients			
Ingredient Name	Strength		
anhydrous citric acid (UNII: XF417D3PSL)			
dextrose (UNII: IY9 XDZ35W2)			
silicon dioxide (UNII: ETJ7Z6XBU4)			
sucralose (UNII: 96K6UQ3ZD4)			

Product Characteristics			
Color	WHITE	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:51596-004-05	5 in 1 BOX			
1	NDC:51596-004-01	1 in 1 POUCH; Type 0: Not a Combination Product			
2	NDC:51596-004-12	12 in 1 BOX			
2	NDC:51596-004-01	1 in 1 POUCH; Type 0: Not a Combination Product			
3	NDC:51596-004-24	24 in 1 BOX			
3	NDC:51596-004-01	1 in 1 POUCH; Type 0: Not a Combination Product			
4	NDC:51596-004-10	10 in 1 BOX			
4		1 in 1 PACKET; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part335	09/01/2011	

Labeler - Breakthrough Products Inc. (962008251)

Revised: 10/2014 Breakthrough Products Inc.