

**MUCUS RELIEF SINUS SEVERE CONGESTION AND PAIN- acetaminophen,  
guaifenesin, and phenylephrine hydrochloride tablet, coated  
CVS PHARMACY**

-----  
**1167-CVS-2020-0908**

***Drug Facts***

<b><i>Active ingredients (in each caplet)</i></b>	<b><i>Purpose</i></b>
Acetaminophen 325 mg	Pain reliever
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves:
  - nasal congestion
  - headache
  - minor aches and pains
  - sinus congestion and pressure
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**When using this product do not use more than directed**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or persistent headache

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Other information**

- store between 20-25°C (68-77°F) in a dry place

- retain outer package for complete product information

### **Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

### **Questions or comments?**

1-844-705-4384

### **PRINCIPAL DISPLAY PANEL**

CVS Health®

Compare to the active ingredients in Mucinex® Sinus-Max® Severe Congestion & Pain\*

NDC 68942-786-02

MAXIMUM STRENGTH

Mucus Relief Sinus Severe Congestion & Pain

ACETAMINOPHEN - Pain reliever

GUAIFENESIN - Expectorant

PHENYLEPHRINE HCl - Nasal decongestant

Relieves:

- Sinus congestion
- Headache

Thins & loosens mucus

20 CAPLETS

Ages 12 Years & Over

Actual Size

**MAXIMUM STRENGTH**  
**Mucus Relief Sinus**  
**Severe Congestion & Pain**



Compare to the active ingredients  
in Mucinex® Sinus-Max®  
Severe Congestion & Pain\*

NDC 69842-786-02

**MAXIMUM STRENGTH**  
**Mucus Relief Sinus**  
**Severe Congestion & Pain**


**ACETAMINOPHEN** - Pain reliever  
**GUAIFENESIN** - Expectorant  
**PHENYLEPHRINE HCl** - Nasal decongestant

**Relieves:**  
• Sinus congestion  
• Headache  
**Thins & loosens mucus**

**20 CAPLETS**

**Ages 12 Years & Over**

Actual Size



**Drug Facts**  
**Active ingredients (in each caplet)**  
Acetaminophen 325 mg.....Pain reliever  
Guaifenesin 200 mg.....Expectorant  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses**  
temporarily relieves:  
■ nasal congestion ■ minor aches and pains  
■ sinus congestion and pressure  
■ promotes nasal and/or sinus drainage  
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
■ more than 12 caplets in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks daily while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
■ diabetes ■ heart disease  
■ liver disease ■ thyroid disease  
■ high blood pressure  
■ trouble urinating due to an enlarged prostate gland  
■ cough that occurs with too much phlegm (mucus)

**Drug Facts (continued)**  
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

**Stop use and ask a doctor if**  
■ nervousness, dizziness, or sleeplessness occur  
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ new symptoms occur  
■ cough comes back or occurs with rash or persistent headache  
These could be signs of a serious condition.  
**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.**  
**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not take more than directed (see Overdose warning)  
■ do not take more than 12 caplets in any 24-hour period  
■ adults and children 12 years and older: take 2 caplets every 4 hours  
■ children under 12 years of age: do not use

**Other information**  
■ store between 20-25°C (68-77°F) in a dry place  
■ retain outer package for complete product information

**Inactive ingredients**  
colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6, aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

\*This product is not manufactured or distributed by Mucinex® Sinus-Max® distributor of Mucinex® Sinus-Max®  
Maximum Strength Per 4-Hour Dose

**Drug Facts (continued)**  
**Questions or comments?**  
1-844-705-4384

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Distributed by: **CVS Pharmacy, Inc.**  
One CVS Drive, Woonsocket, RI 02895  
© 2020 CVS Pharmacy  
CVS.com® 1-800-SHOP CVS  
V-96611  
**CVS Quality**  
Money. Back Guarantee

F116702CVS\_R0

#938777

0 50428 33235 1

# MUCUS RELIEF SINUS SEVERE CONGESTION AND PAIN

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69842-786
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STARCH, PREGELATINIZED CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	AAA;1166
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-786-02	2 in 1 CARTON	09/01/2020	07/31/2026

<b>1</b>	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	09/01/2020	07/31/2026

**Labeler -** CVS PHARMACY (062312574)

Revised: 10/2024

CVS PHARMACY