

ASPIRIN- aspirin tablet, film coated
WinCo Foods, LLC

Winco 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- asthma (wheezing)
- hives
- facial swelling
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor

- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions?

Call **800-824-1706** Monday – Friday 9am – 4pm MST

Principal Display Panel

**Compare to the active ingredient in GENUINE BAYER® Aspirin

WinCo®
FOODS

Aspirin

PAIN RELIEVER - FEVER REDUCER
(NSAID)

300
Tablets

325 mg EACH

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

** This product is not manufactured or distributed by
BAYER AG, distributor of GENUINE BAYER® Aspirin.

100% SATISFACTION GUARANTEE

Comments & Questions? 800-824-1706

50844 REV0122C15717 WF2204

**Compare to the active ingredient in GENUINE BAYER® Aspirin



Drug Facts
Active ingredient (in each tablet) Purpose
 Aspirin 325 mg (NSAID) * Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves:
 ■ minor pain of arthritis
 ■ muscle pain ■ menstrual pain
 ■ headache ■ pain and fever of colds
 ■ toothache

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ asthma (wheezing) ■ hives
 ■ facial swelling ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ take more or for a longer time than directed
 ■ have 3 or more alcoholic drinks every day while using this product

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

** This product is not manufactured or distributed by BAYER AG, distributor of GENUINE BAYER® Aspirin.

100% SATISFACTION GUARANTEE
 Comments & Questions? 800-824-1706



PEEL HERE FOR MORE DRUG FACTS

STOP PEELING

Questions? Call 800-824-1706 Monday - Friday 9am - 4pm MST
Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol

Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Directions
 ■ do not take more than directed
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: do not use unless directed by a doctor

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ ringing in the ears or a loss of hearing occurs
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur ■ redness or swelling is present
 These could be signs of a serious condition.

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis ■ you are taking a diuretic
Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn ■ you have asthma
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you are taking a diuretic

Drug Facts (continued)

WinCo 44-157

ASPIRIN			
aspirin tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-110
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
Inactive Ingredients			
	Ingredient Name	Strength	
	STARCH, CORN (UNII: O8232NY3SJ)		
	HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
	POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)		
	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		

Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;157;ASPIRIN	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-110-30	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
2	NDC:67091-110-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	11/30/2024
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	02/10/1996		

Labeler - WinCo Foods, LLC (056098817)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(67091-110)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(67091-110)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(67091-110)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(67091-110)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(67091-110)