# CHILDRENS PAIN RELIEF- acetaminophen suspension H E B

-----

**Drug Facts** 

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

### **Purpose**

Pain reliever/fever reducer

#### Uses

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
  - the common cold
  - ∘ flu
  - headache
  - sore throat
  - toothache

### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

### Ask a doctor before use if your child has

liver disease.

### Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

### When using this product,

### do not exceed recommended dosage (see overdose warning)

### Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

### Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

#### **Directions**

- this product does not contain directions or complete warnings for adult use
- do not take more than directed (see overdose warning)
- shake well before using
- mL= milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose, use age.

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 ml
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

\*or as directed by a doctor

#### Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- see bottom panel for lot number and expiration date

### Inactive ingredients

acesulfame potassium, butylparaben, carboxymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

### **Questions or Comments?**

Call toll free: 1-877-753-3935 Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

Compare to Children's Tylenol® Oral Suspension active ingredient\*\*

Children's

Pain Relief

Acetaminophen 160 mg per 5 mL

Oral suspension

Pain Reliever/Fever Reducer

For Ages 2 to 11 years

- Alcohol free
- Aspirin free
- Ibuprofen free

Cherry flavor

FL OZ (mL)

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER THE CAP IS BROKEN OR MISSING

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204

#### **Product Label**



#### Pain Reliever/Fever Reducer

#### For Ages 2 to 11 Years

- Alcohol Free
- Aspirin Free
- Ibuprofen Free



#### Drug Facts

#### Active ingredient (in each 5 mL)

Purpose

#### **Uses** temporarily reduces fever

- temporarily relieves minor aches and pains due to ■ the common cold
  ■ flu headache
- sore throat ■ toothache

#### Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist,
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see Overdose warning).

#### Stop use and ask a doctor if new symptoms occur

- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days These could be signs of a serious condition.

#### Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

#### Directions

- m this product does not contain directions or complete warnings for adult use
- do not take more than directed (see Overdose warning)
- shake well before using
- mL = milliliter

#### Drug Facts (continued)

- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (year)	Dose (mL)*	
under 24	under 2	ask a doctor	
24-35	2-3	5 mL	
36-47	4-5	7.5 mL	
48-59	6-8	10 mL	
60-71	9-10	12.5 mL	
72-95	11	15 mL	

\*or as directed by a doctor

#### Other information

- each 5 mL contains: sodium 2 mg store between 20-25°C (68-77°F) do not refrigerate
- see bottom panel for lot number and expiration date

Inactive ingredients acesulfame potassium, butylparaben, carboxymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan

#### Questions or comments?

Call toll free: 1-877-753-3935 Monday-Friday 9AM-5PM EST

"This product is not manufactured or distributed by McNeil Consume Healthcare, distributor of Children's Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

MADE WITH PRIDE AND CARE FOR H-E-B® SAN ANTONIO, TX 78204





PLD-I231D FC008273 8582-1801 Product of China

Lot No.:

Exp. Date:

H-E-B Children's Pain Relief Cherry Flavor

### CHILDRENS PAIN RELIEF

acetaminophen suspension

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-482	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)			
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)			
BUTYLPARABEN (UNII: 3QPI1U3FV8)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERIN (UNII: PDC6A3C0OX)			
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL (UNII: 506T60A25R)			
XANTHAN GUM (UNII: TTV12P4NEE)			

Color Score Shape Size Flavor CHERRY Imprint Code	Product Characteristics			
-				
Flavor CHERRY Imprint Code				
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808- 482-04	1 in 1 BOX	03/08/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/08/2015	

## **Labeler -** H E B (007924756)

Revised: 4/2024 H E B