STOOL SOFTENER DOCUSATE SODIUM- docusate sodium capsule, gelatin coated

Magno-Humphries, Inc.

Stool Softener Docusate Sodium

MHL NDC 54257-902-06

Stool Softener Docusate Sodium 100mg Compare to the active ingredient in Colace *

R

Effective, Gentle Stimulant Free For Comfortable Relief

1000 SOFTGELS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Active ingredient (in each softgel)

Docusate sodium 100mg

Purpose

Stool softener

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 12-72 hours

Warnings

Do not use

Ask a doctor before use if you have

Stop use and ask a doctor if

ask a health professional before use. If pregnant or breast-feeding,

- if you are presently taking mineral oil, unless told by a doctor
- stomach pain
- nausea

- vomiting
- noticed a sudden change in bowel movements that continues over a period of 2 weeks
- you have rectal bleeding or you fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a stool softener for mroe than 1 week.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center () right away. **1-800-222-1222**

Directions

- Take only by mouth. This dose may be taken as a single daily dose or in divided doses.
- take 1-3 softgels daily or as directed by a doctor **adults and children over 12** years of age:
- take 1 softgel daily or as directed by a doctor children 2 to under 12 years of age:
- ask a doctor children under 2 years of age:

Other information

- sodium 7mg each softgel contains:
- store between 15° to 30°C (59° to 86°F)
- Protect from moisture

Inactive ingredients

FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, purified water, sorbitol.

Questions?


call toll-free **1-800-935-6737**

*This product is not manufactured or distributed by the owner of the registered trademark Colace®. Distributed by: Magno-Humphries, Inc. OR 97223 U.S.A. Visit our website: www.magno-humphries.com


trademark Colace®. Distributed by: , OR 97223 U.S.A.

Magno-Humphries, Inc.Visit our website: www.magno-humphries.com

Package Labeling:



1000 SOFTGELS	Effective, Gentle Stimulant Free For Comfortable Relief	Compare to the active ingredient in Colace®*	Docusate Sodium 100 mg	Softener	Stool	4257-902-06		
DO N	ot use if pr	INTED S	EAL UN	DER CAP	IS MISSIN	IG OR	DAMAGE	D
-	Facts						_	
	e ingredie						Purpos	
_	te sodium 10							er
movemen	 relieves occa it in 12-72 hours 	sional cons	tipation (ii	rregularity) •	generally p	roduces	a bowel	
Warn	ings Do not	use • if y	ou are pr	esently takin	g mineral oi	l, unless	told by a do	ctor
Ask a do	ator boforo uno							
		if you have			nausea • v			
 noticed Stop use movement 	a sudden chang and ask a doct after use of a la	e in bowel or if • yo axative. The	movemer u have re ese could	ts that continents that contine ts that contin	nues over a 1 or you fail f	period of to have a	f 2 weeks	
 noticed Stop use movement you need If pregnation Keep out 	a sudden chang and ask a doct	e in bowel or if • yo axative. The softener for ding, ask dren. In ca	movemer u have re ese could more that a health p ase of acc	its that continue that bleeding be signs of a an 1 week. Professional I idental over	nues over a g or you fail t a serious cor before use.	period of to have a ndition.	f 2 weeks a bowel	
noticed Stop use movemen you nee If pregna Keep out Poison Co Direct	a sudden chang and ask a doct at after use of a la d to use a stool nt or breast-fee of reach of chil	e in bowel or if • yo axative. The softener for ding, ask dren. In ca 200-222-12	movemen u have re ese could more that a health p ase of acc 22) right	nts that conti ictal bleeding be signs of a an 1 week. professional l idental overa away.	nues over a j o r you fail f a serious cor before use. dose, get me	period of to have a ndition. edical he	f 2 weeks a bowel Ip or contact	
noticed Stop use movemen you nee If pregna Keep out Poison Co Direct or in divis adults	a sudden chang and ask a doct at after use of a la do to use a stool s nt or breast-fee of reach of chil ntrol Center (1-8 tions • Take	e in bowel or if • yo axative. The softener for ding, ask dren. In ca 000-222-12 only by me	movemen u have re ese could more tha a health p ase of acc 22) right outh. This	nts that conti ctal bleeding be signs of a n 1 week. orofessional l cidental over away.	nues over a j o r you fail f a serious cor before use. dose, get me	period of to have a ndition. edical hel single d	f 2 weeks a bowel Ip or contact laily dose	
noticed Stop use movemer you need If pregna Keep out Poison Co Or in divid adults 12 year • childre	a sudden chang and ask a doct tt after use of a la d to use a stool : nt or breast-fee of reach of chill introl Center (1-8 tions • Take ded doses. and children o	e in bowel or if • yoo axative. The softener for ding, ask dren. In ca 000-222-12 only by mo ver ta	movemer u have re ese could more that a health p ase of acc (22) right outh. This ke 1-3 s	its that conti inctal bleeding be signs of a an 1 week. professional I idental over away. i dose may b oftgels daily	nues over a g or you fail t a serious cor before use. dose, get me e taken as a	period of to have a ndition. edical he single d ted by a	f 2 weeks a bowel Ip or contact laily dose a doctor	
noticed Stop use movemen you nee if pregna Keep out Poison Co Direct or in divi • adults 12 year • childre • childre	a sudden chang and ask a doct tt after use of a la d to use a stool: nt or breast-fee of reach of chil ntrol Center (1-8 tions • Take ded doses. and children o s of age: n 2 to under s of age:	e in bowel pr if • yo axative. The softener for ding, ask dren. In ca 000-222-12 only by mo ver ta ta	movemer u have re ese could more that a health p ase of acc (22) right outh. This ke 1-3 s	Its that conti ctal bleeding be signs of a n 1 week. rofessional li idental over away. close may b oftgels daily or	nues over a j or you fail f a serious cor before use. dose, get me e taken as a r or as direc	period of to have a ndition. edical he single d ted by a	f 2 weeks a bowel Ip or contact laily dose a doctor	
noticed Stop use movemer you need If pregna Keep out Poison Co Or in divits 12 year • childre 12 years Other	a sudden chang and ask a doct it after use of a li d to use a stool : nt or breast-fee of reach of chill introl Center (1-5 tions • Take ded doses. and children o is of age: in 2 to under so f age: an under	e in bowel or if • yo axative. The softener for ding, ask dren. In ca 000-222-12 only by m ver ta ta as 000 • ea	movemer u have re se could more that a health p ase of acc (22) right buth. This ke 1-3 s ke 1 soft sk a doct	its that conti its that conti be signs of a professional li idental over away. e dose may b offgels daily gel daily or or el contains:	nues over a j or you fail ta a serious cor before use. dose, get me e taken as a r or as directed as directed sodium 7 m	period of to have a ndition. edical hei single d ted by a by a do	f 2 weeks a bowel Ip or contact laily dose a doctor	
noticed Stop use movemer you nee If pregna Keep out Poison Co Direct or in divi • adults 12 year • childre 12 year • childre 2 years Other • store be Inacti	a sudden chang and ask a doct it after use of a lik dib use a stool : int or breast-fee of reach of chill introl Center (1-5 tions • Take ded doses. and children o is of age: in 2 to under of age: information	e in bowel or if • yo axative. The softener for ding, ask dren. In c2 000-222-12 only by m ver ta conv • eac C (59° to 84 ients F	movemer u have re- se could more that a health p ase of acc 22) right buth. This ike 1-3 s ike 1 soft sk a doct ch softgr 6°F) • Pr D&C red	Its that conti its that conti tical bleeding be signs of a in 1 week. rofessional idental over away. dose may b offgels daily gel daily or or el contains: otect from m #40, FD&C y	nues over a j or you fail i a serious con before use. Jose, get me e taken as a or as directed as directed sodium 7m obsture	period of to have a adition. edical hel single d ted by a by a do	f 2 weeks a bowel Ip or contact laily dose a doctor cctor	
noticed Stop use movemer you nee If pregna Keep out Poison Co Direct or in divi adults 12 year childre 12 years Other store be Inacti polyethyle	a sudden chang and ask a doct it after use of a lik dit use a stool : nt or breast-fee of reach of chill ntrol Center (1-5 tions • Take ded doses. and children o 's of age: of age: in 2 to under of age: information tween 15 to 30 is ve ingredia	e in bowel or if • yoaxative. The softener for ding, ask dren. In cc woo-222-12 only by mo ver ta ta as OD • eac C (59° to 86 cents F urified wate	movemer u have re use could more that a health pi see of acc 22) right ke 1-3 s ke 1 soft ke 1 soft sk a doct ch softg 3F) • P D&C red	Its that conti its that conti its that conti its that conti be signs of a in 1 week. ordessional idental over away. oftgels daily gel daily or or el contains: rotect from m #40, FD&C y ol.	nues over a j or you fail i a serious con before use. Jose, get me e taken as a or as directed as directed sodium 7m obsture	period of to have a adition. edical hel single d ted by a by a do	f 2 weeks a bowel Ip or contact laily dose a doctor cctor	
noticed Stop use movemer you nee If pregna Keep out Poison Co Direct or in divi adults 12 year childre 12 years othidre torinder store be Inacti polyethyle Quest *This pro	a sudden chang and ask a doct that far use of a la d to use a stool: int or breast-fee of reach of chili ntrol Center (1-8 tions • Take ded doses. and children o s of age: in 2 to under s of age: in a to under s of age: in formatii etween 15 to 30' ive ingredia ene glycol 400, p	e in bowel or if • yoaxative. The softener for ding, ask dren. In c: yoo-222-12 only by mo ver ta ta as OD • eæ C (59' to 80 fents F urified wate oll-free 1- nufactured	movemen u have re esse could a health p more that more that more that more that more that a health p size of acc c22) right that health p size of acc size of acc	Its that conti its that conti its that conti its that conti be signs of a in 1 week. ordessional I idental over away. oftgels daily oftgels daily or or el contains: otect from m #40, FD&C yol. 35-6737 ibuted by	nues over a j or you fail i a serious con before use. Jose, get me e taken as a or as directed as directed sodium 7m obsture	period of to have a adition. edical hel single d ted by a by a do	f 2 weeks a bowel Ip or contact laily dose a doctor cctor	



STOOL SOFTENER DOCUSATE SODIUM

docusate sodium capsule, gelatin coated

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Sou	urce)	NDC:542	57-902
Route of Administration	ORAL				
Active Ingredient/Active	Moiety				
Ingredient Name Basis of Strength Stren					Strength

100 mg

Inactive Ingre	edients					
	Strength					
GELATIN, UNSPEC	IFIED (UNII:	2G86QN327L)				
GLYCERIN (UNII: PE	DC6A3C0OX)					
POLYETHYLENE G	LYCOL 400	(UNII: B697894	SGQ)			
WATER (UNII: 059Q	F0KO0R)					
SORBITOL (UNII: 5	06T60A25R)					
Product Chara	acteristic	S				
		red	Score		no score	
Shape		OVAL	Size		12mm	
Flavor			Imprint Code		SCU1	
Contains						
Packaging						
# Item Code		Package Description		Marketing Start Date	Marketing End Date	
1 NDC:54257-902- 06	1000 in 1 E Product	OTTLE; Type 0:	Not a Combination	04/04/2017		
Markating	Inform	otion				
Marketing						
Mayleting	Appl	Application Number or Monograph Citation		Marketing Star Date	t Marketing End Date	
Marketing Category		Cita		2410	Butt	

Labeler - Magno-Humphries, Inc. (063251433)

Revised: 10/2023

Magno-Humphries, Inc.