MINI NICOTINE- nicotine polacrilex lozenge ATLANTIC BIOLOGICALS CORP.

Drug Facts

Active ingredient (in each lozenge)

Nicotine polacrilex 4 mg

Purpose

Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding,

only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

diarrhea, weakness and rapid heartbeat

• you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions (4 mg lozenge)

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2	1 lozenge every 2 to 4	1 lozenge every 4 to 8
hours	hours	hours

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information

- Phenylketonurics: Contains Phenylalanine 2.8 mg per lozenge
- store at 20-25°C (68-77°F)
- keep vial tightly closed and protect from light

Inactive ingredients

aspartame, calcium polycarbophil, flavor, maltodextrin, mannitol, sodium alginate, sodium bicarbonate, sodium stearyl fumarate, talc, xanthan gum

Questions or comments?

Call **1-800-645-2158**

DISTRIBUTED BY:

- ATLANTIC BIOLOGICALS CORP.
- MIAMI, FL 33179

Package Label (4 mg lozenge)

17856-1241-01 NICOTINE MINI (POLACRILEX) DELIVERS 1LOZ/4mg STOP SMOKING AID

See package insert for indications and dosage schedule

For individuals who smoke their first cigarette within 30 minutes of waking up. Contains 2.8mg of Phenylalanine per lozonge. Sugar Free Mint Flavor. Keep tighly closed and protect from light. Store at 20°-25°C(68°-77°F). **Keep this and all Medications out of the reach of children**



Distributed by: Atlantic Biologicals Corp. Miami, FL 33179

Ð

UDDSE

2

Rev.08/21 Call to Reorder: 800.509.7592

MINI NICOTINE nicotine polacrilex lozenge			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:17856-1241(NDC:0536-1241)
Route of Administration	ORAL		

	ctive Ingredi	ent/Active	Moiety						
Ingredient Name				Basis of Strength		Strength			
NI	COTINE (UNII: 6M	13C89ZY6R) (NIC	COTINE - UNII:6	M3C89ZY6R)		NICOTINE		4 mg	
In	active Ingre	dients							
			Ingredient	Name			S	Strength	
AS	PARTAME (UNII:	Z0H242BBR1)							
CA		RBOPHIL (UNII:	8F049NKY49)						
M	ALTODEXTRIN (U	NII: 7CVR7L4A2I	D)						
M	ANNITOL (UNII: 30	OWL53L36A)							
sc	DIUM ALGINATE	(UNII: C269C40	G2ZQ)						
50	DIUM BICARBO	NATE (UNII: 8MI	DF5V39QO)						
5C	DIUM STEARYL	FUMARATE (UI	NII: 7CV7WJK4U	I)					
ТА	LC (UNII: 7SEV7J4	IR1U)							
XA	NTHAN GUM (UN	III: TTV12P4NEE)						
Pı	roduct Chara	octeristics							
Co	olor	WHI	TE	Score			no score		
Sł	аре	OVA	AL.	Size			10mm		
Fla	avor	MIN	Т	Imprint Code		131			
Contains									
Pa	ackaging								
#	ltem Code	Рас	kage Descr	iption	Marl	ceting Start Date		eting End Date	
1	NDC:17856- 1241-1	50 in 1 BOX		05/08/2024					
1	NDC:17856- 1241-2	1 in 1 POUCH; Type 0: Not a Combination Product							
Μ	arketing	Informati	ion						
	Marketing	Applicat	ion Number	or Monograph	n M	arketing Start	Mar	keting End	
	Category		Citatio	n		Date		Date	

Labeler - ATLANTIC BIOLOGICALS CORP. (047437707)

Registrant - ATLANTIC BIOLOGICALS CORP. (047437707)

Establishment						
Name	Address	ID/FEI	Business Operations			
UNIT DOSE SOLUTIONS		360804194	repack(17856-1241)			