

**KROGER DAYTIME AND NIGHTTIME COLD AND FLU SOFTGEL- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, doxylamine succinate**

**The Kroger Co.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----

**Kroger DayTime and NightTime Cold & Flu SOFTGELS**

**DayTime Cold & Flu**

***Drug Facts***

***Active ingredients (in each softgel)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Nasal decongestant

***Uses***

temporarily relieves common cold/flu symptoms:

• nasal congestion • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 softgels in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease • heart disease
- high blood pressure
- thyroid disease • diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**When using this product, do not use more than directed****Stop use and ask a doctor if** • you get nervous, dizzy, or sleepless

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present • new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- store at room temperature between 20°-25°C (68°-77°F)
- protect from light, heat, and moisture

### ***Inactive ingredients***

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, sorbitol sorbitan anhydrous, titanium dioxide

### ***Questions or comments?***

**1-800-632-6900**

### **NightTime Cold & Flu**

#### ***Drug Facts***

#### ***Active ingredients (in each softgel)***

Acetaminophen 325 mg  
Dextromethorphan HBr 15 mg  
Doxylamine succinate 6.25 mg

#### ***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Antihistamine

#### ***Uses***

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

#### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 softgels in 24 hrs, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### ***Do not use***

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease • heart disease • thyroid disease • diabetes
- glaucoma • cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers • taking the blood thinning drug warfarin

**When using this product • do not use more than directed**

- excitability may occur, especially in children
- marked drowsiness may occur • avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present
- new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

***Other information***

- store at room temperature between 20°-25°C (68°-77°F)
- protect from light, heat, and moisture

***Inactive ingredients***

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol 400, polyvinyl

pyrrolidone, propylene glycol, sorbitol sorbitan anhydrous, titanium dioxide

***Questions or comments?***

**1-800-632-6900**

**COMPARE TO active ingredients of VICKS® DAYQUIL®/DAYQUIL® COLD & FLU RELIEF LIQUICAPS™\***

**POWERFUL RELIEF**

DAYTIME OR NIGHTTIME

Multi-Symptom Relief

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**DISTRIBUTED BY THE KROGER CO.**

**CINCINNATI, OHIO**

**QUALITY GUARANTEE**

**[www.kroger.com](http://www.kroger.com)**

**PRODUCT OF INDIA**

\*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the trademarks Vicks®, DayQuil™, NyQuil™ and LiquiCaps™

REV.00-092024

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**Packaging**

## DRUG FACTS LABEL

**COMPARE TO active ingredients of VICKS® DAYQUIL®/DAYQUIL®  
COLD & FLU RELIEF LIQUICAPS™ \***

**POWERFUL RELIEF**  
DAYTIME OR NIGHTTIME

**Cold & Flu**


**DayTime**

Multi-Symptom Relief

**Acetaminophen**  
Dextromethorphan HBr  
Phenylephrine HCl

Pain Reliever  
Fever Reducer  
Cough Suppressant  
Nasal Decongestant

**ACTUAL SIZE**



**32 SOFTGELS**

**Cold & Flu**


**NightTime**

Multi-Symptom Relief

**Acetaminophen**  
Dextromethorphan HBr  
Doxylamine Succinate

Pain Reliever  
Fever Reducer  
Cough Suppressant  
Antihistamine

**ACTUAL SIZE**



**16 SOFTGELS**

[illegible]

Drug Facts (continued)		Drug Facts (continued)	
<p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• with any other drug containing acetaminophen (prescription or over-the-counter).</li> <li>• if you are now taking a prescription narcotic or other substance that has a danger of respiratory depression or other central nervous system (CNS) depression.</li> <li>• if you are now taking a prescription narcotic or other substance that has a danger of respiratory depression or other central nervous system (CNS) depression.</li> <li>• if you are now taking a prescription narcotic or other substance that has a danger of respiratory depression or other central nervous system (CNS) depression.</li> </ul>		<p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• with any other drug containing acetaminophen (prescription or over-the-counter).</li> <li>• if you are now taking a prescription narcotic or other substance that has a danger of respiratory depression or other central nervous system (CNS) depression.</li> <li>• if you are now taking a prescription narcotic or other substance that has a danger of respiratory depression or other central nervous system (CNS) depression.</li> <li>• if you are now taking a prescription narcotic or other substance that has a danger of respiratory depression or other central nervous system (CNS) depression.</li> </ul>	
<p><b>Ask a doctor before using this product:</b></p> <ul style="list-style-type: none"> <li>• if you have a history of asthma, chronic bronchitis, or emphysema.</li> <li>• if you have a history of asthma, chronic bronchitis, or emphysema.</li> <li>• if you have a history of asthma, chronic bronchitis, or emphysema.</li> </ul>		<p><b>Ask a doctor before using this product:</b></p> <ul style="list-style-type: none"> <li>• if you have a history of asthma, chronic bronchitis, or emphysema.</li> <li>• if you have a history of asthma, chronic bronchitis, or emphysema.</li> <li>• if you have a history of asthma, chronic bronchitis, or emphysema.</li> </ul>	
<p><b>Warnings:</b></p> <ul style="list-style-type: none"> <li>• <b>Do not take more than 4 tablets in 24 hours.</b></li> <li>• <b>Do not take more than 4 tablets in 24 hours.</b></li> <li>• <b>Do not take more than 4 tablets in 24 hours.</b></li> </ul>		<p><b>Warnings:</b></p> <ul style="list-style-type: none"> <li>• <b>Do not take more than 4 tablets in 24 hours.</b></li> <li>• <b>Do not take more than 4 tablets in 24 hours.</b></li> <li>• <b>Do not take more than 4 tablets in 24 hours.</b></li> </ul>	
<p><b>Directions:</b></p> <ul style="list-style-type: none"> <li>• <b>Adults and children 12 years &amp; over:</b> 2 softgels with water every 4 hrs.</li> <li>• <b>Children 6 to 11 years:</b> 1 softgel with water every 4 hrs.</li> <li>• <b>Children 4 to 5 years:</b> 1 softgel with water every 4 hrs.</li> </ul>		<p><b>Directions:</b></p> <ul style="list-style-type: none"> <li>• <b>Adults and children 12 years &amp; over:</b> 2 softgels with water every 6 hrs.</li> <li>• <b>Children 6 to 11 years:</b> 1 softgel with water every 6 hrs.</li> <li>• <b>Children 4 to 5 years:</b> 1 softgel with water every 6 hrs.</li> </ul>	
<p><b>Other information:</b></p> <ul style="list-style-type: none"> <li>• <b>Store at room temperature between 20°-25°C (68°-77°F).</b></li> <li>• <b>Protect from light, heat, and moisture.</b></li> <li>• <b>Protect from light, heat, and moisture.</b></li> </ul>		<p><b>Other information:</b></p> <ul style="list-style-type: none"> <li>• <b>Store at room temperature between 20°-25°C (68°-77°F).</b></li> <li>• <b>Protect from light, heat, and moisture.</b></li> <li>• <b>Protect from light, heat, and moisture.</b></li> </ul>	
<p><b>Inactive ingredients:</b></p> <ul style="list-style-type: none"> <li>• <b>FD-302, FD-303, FD-304, FD-305, FD-306, FD-307, FD-308, FD-309, FD-310, FD-311, FD-312, FD-313, FD-314, FD-315, FD-316, FD-317, FD-318, FD-319, FD-320, FD-321, FD-322, FD-323, FD-324, FD-325, FD-326, FD-327, FD-328, FD-329, FD-330, FD-331, FD-332, FD-333, FD-334, FD-335, FD-336, FD-337, FD-338, FD-339, FD-340, FD-341, FD-342, FD-343, FD-344, FD-345, FD-346, FD-347, FD-348, FD-349, FD-350, FD-351, FD-352, FD-353, FD-354, FD-355, FD-356, FD-357, FD-358, FD-359, FD-360, FD-361, FD-362, FD-363, FD-364, FD-365, FD-366, FD-367, FD-368, FD-369, FD-370, FD-371, FD-372, FD-373, FD-374, FD-375, FD-376, FD-377, FD-378, FD-379, FD-380, FD-381, FD-382, FD-383, FD-384, FD-385, FD-386, FD-387, FD-388, FD-389, FD-390, FD-391, FD-392, FD-393, FD-394, FD-395, FD-396, FD-397, FD-398, FD-399, FD-400, FD-401, FD-402, FD-403, FD-404, FD-405, FD-406, FD-407, FD-408, FD-409, FD-410, FD-411, FD-412, FD-413, FD-414, FD-415, FD-416, FD-417, FD-418, FD-419, FD-420, FD-421, FD-422, FD-423, FD-424, FD-425, FD-426, FD-427, FD-428, FD-429, FD-430, FD-431, FD-432, FD-433, FD-434, FD-435, FD-436, FD-437, FD-438, FD-439, FD-440, FD-441, FD-442, FD-443, FD-444, FD-445, FD-446, FD-447, FD-448, FD-449, FD-450, FD-451, FD-452, FD-453, FD-454, FD-455, FD-456, FD-457, FD-458, FD-459, FD-460, FD-461, FD-462, FD-463, FD-464, FD-465, FD-466, FD-467, FD-468, FD-469, FD-470, FD-471, FD-472, FD-473, FD-474, FD-475, FD-476, FD-477, FD-478, FD-479, FD-480, FD-481, FD-482, FD-483, FD-484, FD-485, FD-486, FD-487, FD-488, FD-489, FD-490, FD-491, FD-492, FD-493, FD-494, FD-495, FD-496, FD-497, FD-498, FD-499, FD-500, FD-501, FD-502, FD-503, FD-504, FD-505, FD-506, FD-507, FD-508, FD-509, FD-510, FD-511, FD-512, FD-513, FD-514, FD-515, FD-516, FD-517, FD-518, FD-519, FD-520, FD-521, FD-522, FD-523, FD-524, FD-525, FD-526, FD-527, FD-528, FD-529, FD-530, FD-531, FD-532, FD-533, FD-534, FD-535, FD-536, FD-537, FD-538, FD-539, FD-540, FD-541, FD-542, FD-543, FD-544, FD-545, FD-546, FD-547, FD-548, FD-549, FD-550, FD-551, FD-552, FD-553, FD-554, FD-555, FD-556, FD-557, FD-558, FD-559, FD-560, FD-561, FD-562, FD-563, FD-564, FD-565, FD-566, FD-567, FD-568, FD-569, FD-570, FD-571, FD-572, FD-573, FD-574, FD-575, FD-576, FD-577, FD-578, FD-579, FD-580, FD-581, FD-582, FD-583, FD-584, FD-585, FD-586, FD-587, FD-588, FD-589, FD-590, FD-591, FD-592, FD-593, FD-594, FD-595, FD-596, FD-597, FD-598, FD-599, FD-600, FD-601, FD-602, FD-603, FD-604, FD-605, FD-606, FD-607, FD-608, FD-609, FD-610, FD-611, FD-612, FD-613, FD-614, FD-615, FD-616, FD-617, FD-618, FD-619, FD-620, FD-621, FD-622, FD-623, FD-624, FD-625, FD-626, FD-627, FD-628, FD-629, FD-630, FD-631, FD-632, FD-633, FD-634, FD-635, FD-636, FD-637, FD-638, FD-639, FD-640, FD-641, FD-642, FD-643, FD-644, FD-645, FD-646, FD-647, FD-648, FD-649, FD-650, FD-651, FD-652, FD-653, FD-654, FD-655, FD-656, FD-657, FD-658, FD-659, FD-660, FD-661, FD-662, FD-663, FD-664, FD-665, FD-666, FD-667, FD-668, FD-669, FD-670, FD-671, FD-672, FD-673, FD-674, FD-675, FD-676, FD-677, FD-678, FD-679, FD-680, FD-681, FD-682, FD-683, FD-684, FD-685, FD-686, FD-687, FD-688, FD-689, FD-690, FD-691, FD-692, FD-693, FD-694, FD-695, FD-696, FD-697, FD-698, FD-699, FD-700, FD-701, FD-702, FD-703, FD-704, FD-705, FD-706, FD-707, FD-708, FD-709, FD-710, FD-711, FD-712, FD-713, FD-714, FD-715, FD-716, FD-717, FD-718, FD-719, FD-720, FD-721, FD-722, FD-723, FD-724, FD-725, FD-726, FD-727, FD-728, FD-729, FD-730, FD-731, FD-732, FD-733, FD-734, FD-735, FD-736, FD-737, FD-738, FD-739, FD-740, FD-741, FD-742, FD-743, FD-744, FD-745, FD-746, FD-747, FD-748, FD-749, FD-750, FD-751, FD-752, FD-753, FD-754, FD-755, FD-756, FD-757, FD-758, FD-759, FD-760, FD</b></li></ul>			

READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION

**FPO  
UPC**

NO VARNISH

LOT:  
EXP:

NO COATING  
AREANO COATING  
AREA



## Drug Facts (Nighttime)

### Active ingredients (in each softgel) Purpose

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 15 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine

**Uses** temporarily relieves common cold/flu symptoms:  
■ cough due to minor throat & bronchial irritation ■ sore throat  
■ headache ■ minor aches & pains ■ fever ■ runny nose & sneezing

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 softgels in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Drug Facts (Daytime)

### Active ingredients (in each softgel) Purpose

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:  
■ nasal congestion ■ cough due to minor throat & bronchial irritation  
■ sore throat ■ headache ■ minor aches & pains ■ fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 softgels in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Drug Facts (continued)

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

■ liver disease ■ heart disease ■ thyroid disease ■ diabetes  
■ glaucoma ■ cough that occurs with too much phlegm (mucus)  
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema  
■ trouble urinating due to enlarged prostate gland

### Ask a doctor or pharmacist before use if you are

■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

### When using this product ■ do not use more than directed

■ excitability may occur, especially in children  
■ marked drowsiness may occur ■ avoid alcoholic drinks  
■ be careful when driving a motor vehicle or operating machinery  
■ alcohol, sedatives, and tranquilizers may increase drowsiness

### Stop use and ask a doctor if

■ pain, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present  
■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Drug Facts (continued)

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

■ liver disease ■ heart disease  
■ high blood pressure  
■ thyroid disease ■ diabetes  
■ trouble urinating due to enlarged prostate gland  
■ cough that occurs with too much phlegm (mucus)  
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

### Ask a doctor or pharmacist before use if you are

■ taking the blood thinning drug warfarin

### When using this product, do not use more than directed

**Stop use and ask a doctor if** ■ you get nervous, dizzy, or sleepless  
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Drug Facts (continued)

### Directions

■ take only as directed  
■ do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over 2 softgels with water every 6 hrs

## Drug Facts (continued)

### Directions

■ take only as directed  
■ do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over 2 softgels with water every 4 hrs

children 4 to under 12 yrs	ask a doctor	children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use	children under 4 yrs	do not use
<b>Other information</b> <ul style="list-style-type: none"> <li>■ store at room temperature between 20°-25°C (68°-77°F)</li> <li>■ protect from light, heat, and moisture</li> </ul>		<b>Other information</b> <ul style="list-style-type: none"> <li>■ store at room temperature between 20°-25°C (68°-77°F)</li> <li>■ protect from light, heat, and moisture</li> </ul>	
<b>Inactive ingredients</b> D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol 400, polyvinyl pyrrolidone, propylene glycol, sorbitol sorbitan anhydrous, titanium dioxide		<b>Inactive ingredients</b> FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, sorbitol sorbitan anhydrous, titanium dioxide	
<b>Questions or comments? 1-800-632-6900</b>		<b>Questions or comments? 1-800-632-6900</b>	

## KROGER DAYTIME AND NIGHTTIME COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, doxylamine succinate kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59450-048
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59450-048-48	1 in 1 CARTON; Type 0: Not a Combination Product	01/08/2025	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
<b>Part 1</b>	4 BLISTER PACK	32
<b>Part 2</b>	4 BLISTER PACK	16

## Part 1 of 2

## KROGER DAYTIME COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

### Product Information

<b>Item Code (Source)</b>	NDC:59450-046
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety



Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	A09
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/08/2025	

### Part 2 of 2

### KROGER NIGHTTIME COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

### Product Information

Item Code (Source)	NDC:59450-047			
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	15 mg	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)		DOXYLAMINE SUCCINATE	6.25 mg	
Inactive Ingredients				
Ingredient Name			Strength	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)				
GLYCERIN (UNII: PDC6A3C0OX)				
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SORBITOL (UNII: 506T60A25R)				
SORBITAN (UNII: 6O92ICV9RU)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	green	Score	no score	
Shape	OVAL	Size	19mm	
Flavor		Imprint Code	902	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012		01/08/2025	
Marketing Information				

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/08/2025	

**Labeler -** The Kroger Co. (006999528)

Revised: 1/2025

The Kroger Co.