

IBUPROFEN- ibuprofen capsule, liquid filled
MEIJER DISTRIBUTION INC.

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause delivery problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

Inactive ingredients

FD&C Blue #1, gelatin, glycerin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

For India Manufacturing Site:

FD&C Blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

Questions or comments?

1-888-235-2466

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH “ SEALED for YOUR PROTECTION” IS BROKEN OR MISSING.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

OUR QUALITY GUARANTEE

www.meijer.com/Satisfaction

DIST. MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

L0000845

R0424

For India Manufacturing Site:

P3289/0025 **MADE IN INDIA**

Code: TN/DRUGS/TN/25/00074

R0125 L0000984

20's Carton

meijer

NDC 41250-292-11

COMPARE TO THE

ACTIVE INGREDIENT

IN ADVIL® LIQUI-GELS®†

IbuprofeN

MINI

CAPSULES

Capsules 200 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

SMALLER CAPSULE

SAME STRENGTH

20 Softgels**

****Liquid-Filled Capsules**

meijer.
ibuprofen
Capsules | 200 mg

COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS®†

NDC 41250-292-11

MINI CAPSULES

PAIN RELIEVER/FEVER REDUCER (NSAID) SMALLER CAPSULE SAME STRENGTH**

20 Softgels**

Actual Size

Drug Facts (continued)

Other information: Each capsule contains: potassium 20 mg. Read all warnings and directions before use. Keep carton. Store at 20° to 25°C (68° to 77°F). Avoid excessive heat above 40°C (104°F). Protect from light.

Inactive ingredients: FD&C Blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol, and sorbitol.

Questions or comments? Call 1-888-235-2466

† Do not use if you are allergic to any of the ingredients listed above. For a complete list of ingredients, see the back of the carton.

**** 1 liquid-filled capsule is equivalent to 200 mg ibuprofen (NSAID)†.**

Dist. Meijer Distribution, Inc.
Grand Rapids, MI 49544
www.meijer.com
PID 4967780

OUR QUALITY GUARANTEE
The Superior Quality

7 13733 37743 0

Lot No.:
Exp. Date:

Drug Facts (continued)

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - heart burn
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part of your body
 - slurred speech
 - leg swelling
- your gas worsens or lasts more than 10 days
- your gas worsens or lasts more than 3 days
- redness or swelling is present in the genital area
- any new symptoms appear

pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Drug Facts (continued)

Use: Take with food or milk if stomach upset occurs.

When using this product:

- take with food or milk if stomach upset occurs
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
- do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer (right before or after heart surgery)
- ask a doctor before use if you have stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

Warnings:

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach pain
- skin rash/itching
- stomach bleeding
- stomach pain
- stomach bleeding
- stomach pain
- stomach bleeding

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), or have a history of stomach ulcers or bleeding.

Caution: Do not take more than directed.

Drug Facts (continued)

Active ingredient (in each capsule): Solubilized ibuprofen equivalent to 200 mg ibuprofen (NSAID)†.† nonsteroidal anti-inflammatory drug (present as the free acid and potassium salt).

Purpose: Pain reliever/fever reducer.

Uses: Temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscle aches
- minor pain of arthritis
- temporarily reduces fever

80's carton- Made In India label
meijer
NDC 41250-292-78
COMPARE TO THE
ACTIVE INGREDIENT
IN ADVIL® LIQUI-GELS®†
Ibuprofen
MINI
CAPSULES
Capsules 200 mg
PAIN RELIEVER/FEVER REDUCER (NSAID)


SMALLER CAPSULE

SAME STRENGTH

80 Softgels**

****Liquid-Filled Capsules**

(MADE IN INDIA)



IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:41250-292

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	blue	Score	no score
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	IB4
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-292-11	1 in 1 CARTON	03/19/2020	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-292-78	1 in 1 CARTON	03/19/2020	
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-292-83	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/19/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	03/19/2020	

Labeler - MEIJER DISTRIBUTION INC. (006959555)

Registrant - Bionpharma Inc. (079637826)

Establishment

Name	Address	ID/FEI	Business Operations
Patheon Softgels Inc.		002193829	manufacture(41250-292)

Establishment

Name	Address	ID/FEI	Business Operations
ESJAY PHARMA PRIVATE LIMITED		957309497	manufacture(41250-292)

Revised: 4/2025

MEIJER DISTRIBUTION INC.