

PROCTOZONE GMAX- glycerin suppository
Rising Pharma Holdings, Inc.

Proctozone-GMax™ Glycerin Suppositories 2 g

Drug Facts

Active ingredient (in each suppository)

Glycerin USP, 2 g

Purpose

Laxative

Uses

- For relief of occasional constipation
- This product usually produces bowel movement in 1/4 to 1 hour

Warnings

For rectal use only

Do not use for longer than 1 week unless directed by a doctor.

Ask a doctor before use if you have

- Abdominal pain, nausea or vomiting
- A sudden change in bowel habits lasting more than 2 weeks

When using this product it may cause rectal discomfort or a burning sensation

Stop use and ask a doctor if you have

- Rectal bleeding
- No bowel movement after using this product
These may indicate a serious condition.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Tear off one suppository from the strip
- If the suppository seems soft, place it in the refrigerator before use or run cold water over it before removing the foil wrapper
- Gently remove suppository by peeling apart the wrapper (**see illustration 1**)
- Rub a water-based lubricant over the tip or dip it in water. It will help you slide it in smoothly
- Lie on your side and insert 1 suppository, tapered end first, about the length of your finger into the rectum (**see illustration 2**)
- Hold the suppository for about 15-20 minutes to allow time for it to melt
- Adults and children 6 years of age and over..... 1 suppository once daily
- Children under 6 years age..... do not use

Other Information

Keep away from excessive heat.

Inactive ingredients

purified water, sodium stearate.

Questions or Comments:

call 1-844-474-7464

TAMPER EVIDENT, INDIVIDUALLY SEALED FOR YOUR PROTECTION

Distributed by:

Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

Made in India

Code No.: MH/DRUGS/25-KD/724

Issued: 05/2025

NDC 57237-339-21 (2 blisters of 6 suppositories)

Proctozone-GMax™

GLYCERIN SUPPOSITORIES 2 g

ADULT'S LAXATIVE

For Gentle, Predictable Relief of Occasional Constipation

GTM No. 00357237338213
SR. No.: XXXXXXXXXX
LOT: XXXXXXXXX
EXP: YYYYYMM



<p>Drug Facts Active ingredient (in each suppository) Glycerin USP, 2 g Laxative</p>		<p>Warnings For rectal use only. Do not use for longer than 1 week unless directed by a doctor. Ask a doctor before use if you have: ■ A sudden change in bowel habits lasting more than 2 weeks ■ Abdominal pain, nausea or vomiting When using this product it may cause rectal discomfort or a burning sensation. Stop use and ask a doctor if you have: ■ Rectal bleeding ■ No bowel movement after using the product. These may indicate a serious condition. Keep out of reach of children If swallowed, get medical help or contact a Poison Control Center right away.</p>
<p>Directions Tear off one suppository from the strip. If the suppository seems soft, place it in the refrigerator before use or run cold water over it before removing the foil wrapper. Gently remove suppository by peeling apart the wrapper (see illustration 1). Rub a water-based lubricant over the tip or dip it in water. It will help you slide it in smoothly. Lie on your side and insert 1 suppository, tapered end first, about the length of your finger into the rectum (see illustration 2). Hold the suppository for about 15 - 20 minutes to allow time for it to melt. Adults and children 6 years of age and over: 1 suppository once daily. Children under 6 years: do not use.</p>		
<p>Other Information Keep away from excessive heat.</p>		<p>Questions or Comments: call 1-844-474-7464</p>
<p>Inactive ingredients purified water, sodium stearate.</p>		

UNVARNISHED AREA FOR OPZ



1805




Front Foil (57237-339-16)

Annotation: Lot No. and Exp. Date will be embossed on each cavity on back foil as follows:

L.XXXXXXX

E.YYYY/MM



Back Foil (57237-339-16)

Print Repeat 68,55 mm

22.85 mm
(One eye mark distance)

18	26	16	18	26	16
<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>			<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>		
<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>			<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>		
<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>			<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>		
<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>			<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>		
<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>			<p>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</p>		

120.00 mm

NDC 57237-339-52 (5 blisters of 5 suppositories)

GTIN No. 00357237339527
SR. No.: XXXXXXXXXXXXXXXXXX
LOT: XXXXXXXX
EXP: YYYY/MM




UNVARNISHED AREA FOR OPZ

<p>Drug Facts Active ingredient (in each suppository) Glycerin USP, 2 g Purpose Laxative</p>		<p>Uses For relief of occasional constipation This product usually produces bowel movement in 1/4 to 1 hour</p>	
<p>Warnings For rectal use only Do not use for longer than 1 week unless directed by a doctor.</p>		<p>Ask a doctor before use if you have Abdominal pain, nausea or vomiting A sudden change in bowel habits lasting more than 2 weeks When using this product it may cause rectal discomfort or a burning sensation</p>	
<p>Stop use and ask a doctor if you have Rectal bleeding No bowel movement after using this product These may indicate a serious condition.</p>		<p>Keep out of reach of children If swallowed, get medical help or contact a Poison Control Center right away.</p>	
<p>Directions Tear off one suppository from the strip If the suppository seems soft, place it in the refrigerator before use or run cold water over it before removing the foil wrapper Gently remove suppository by peeling apart the wrapper (see illustration 1) Rub a water-based lubricant over the tip or dip it in water. It will help you slide it in smoothly Lie on your side and insert 1 suppository, tapered end first, about the length of your finger into the rectum (see illustration 2) Hold the suppository for about 15 - 20 minutes to allow time for it to melt</p>		<p>Other Information Keep away from excessive heat. Children under 6 years do not use</p>	
<p>Drug Facts (Continued)</p>		<p>Questions or Comments: Call 1-844-474-7464</p>	



Front Foil (57237-339-51)

Annotation: Lot No. and Exp. Date will be embossed on each cavity on back foil as follows:
 L.XXXXXXX
 E.YYYY/MM



Back Foil (57237-339-51)



PROCTOZONE GMAX

glycerin suppository

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57237-339
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	2 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	

Product Characteristics

Color	white (clear)	Score	
Shape	BULLET	Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57237-339-21	2 in 1 CARTON	07/31/2025	
1	NDC:57237-339-16	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:57237-339-52	5 in 1 CARTON	07/31/2025	
2	NDC:57237-339-51	5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007.54	07/31/2025	

Labeler - Rising Pharma Holdings, Inc. (116880195)

Establishment

Name	Address	ID/FEI	Business Operations
Bliss GVS Pharma Limited		675630493	manufacture(57237-339) , analysis(57237-339) , pack(57237-339)

Revised: 1/2026

Rising Pharma Holdings, Inc.