

PROCTOZONE GMAX- glycerin suppository
Rising Pharma Holdings, Inc.

Proctozone-GMax™ Glycerin Suppositories 2 g

Drug Facts

Active ingredient (in each suppository)

Glycerin USP, 2 g

Purpose

Laxative

Uses

- For relief of occasional constipation
- This product usually produces bowel movement in 1/4 to 1 hour

Warnings

For rectal use only

Do not use for longer than 1 week unless directed by a doctor.

Ask a doctor before use if you have

- Abdominal pain, nausea or vomiting
- A sudden change in bowel habits lasting more than 2 weeks

When using this product it may cause rectal discomfort or a burning sensation

Stop use and ask a doctor if you have

- Rectal bleeding
 - No bowel movement after using this product
- These may indicate a serious condition.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Tear off one suppository from the strip
- If the suppository seems soft, place it in the refrigerator before use or run cold water over it before removing the foil wrapper
- Gently remove suppository by peeling apart the wrapper **(see illustration 1)**
- Rub a water-based lubricant over the tip or dip it in water. It will help you slide it in smoothly
- Lie on your side and insert 1 suppository, tapered end first, about the length of your finger into the rectum **(see illustration 2)**
- Hold the suppository for about 15-20 minutes to allow time for it to melt
- Adults and children 6 years of age and over..... 1 suppository once daily
- Children under 6 years age..... do not use



Other Information

Drug Facts Active ingredient (in each suppository) _____ Purpose _____ Glycerin USP, 2 g. _____ Laxative _____		Drug Facts Uses ■ For relief of occasional constipation ■ This product usually produces bowel movement in 1/4 to 1 hour		Warnings For rectal use only Do not use for longer than 1 week unless directed by a doctor. Ask a doctor before use if you have ■ Abdominal pain, nausea or vomiting ■ A sudden change in bowel habits lasting more than 2 weeks When using this product it may cause rectal discomfort or a burning sensation Stop use and ask a doctor if you have ■ Rectal bleeding ■ No bowel movement after using this product These may indicate a serious condition. Keep out of reach of children If swallowed, get medical help or contact a Poison Control	
Questions or Comments: call 1-844-474-7464		Other Information Keep away from excessive heat. Inactive ingredients purified water, sodium stearate.		Directions 1. If the suppository seems soft, place it in the refrigerator before use or run cold water over it before removing the foil wrapper. 2. Gently remove suppository by peeling apart the wrapper (see illustration 1). 3. Rub a water-based lubricant over the tip or dip it in water. It will help you slide it in smoothly. 4. Lie on your side and insert 1 suppository, tapered end first, about the length of your finger into the rectum (see illustration 2). 5. Hold the suppository for about 15 - 20 minutes to allow time for it to melt. Adults and children 6 years of age and over: 1 suppository once daily Children under 6 years: do not use	



Front Foil (57237-339-16)

Annotation: Lot No. and Exp. Date will be embossed on each cavity on back foil as follows:
L.XXXXXXX
E.YYYY/MM



BATCH DETAILS WILL BE EMBOSSED HERE

Back Foil (57237-339-16)

Print Repeat 68,55 mm

22,85 mm
(One eye mark distance)

18	26	16	18	26	16
<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>			<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>		
<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>			<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>		
<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>			<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>		
<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>			<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>		
<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>			<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>		
<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>			<div><div>Diet. By: Riesing Pharma Holdings, Inc. Store at 20-25°C (68-77°F)</div><div><div>N</div><div>3</div><div>57237*33916</div><div>9</div></div></div>		

120.00 mm

NDC 57237-339-52 (5 blisters of 5 suppositories)

GTIN No.00357237339527
SR. No.: XXXXXXXXXX
LOT: XXXXXXXX
EXP: YYYY/MM



UNVARNISHED AREA FOR OPZ

Drug Facts

Active ingredient (in each suppository)
Glycerin USP, 2 g
Purpose
Laxative

Uses

■ For relief of occasional constipation
■ This product usually produces bowel movement in 1/4 to 1 hour

Warnings

For rectal use only
Do not use for longer than 1 week unless directed by a doctor.
Ask a doctor before use if you have
■ Abdominal pain, nausea or vomiting
■ A sudden change in bowel habits lasting more than 2 weeks
When using this product it may cause rectal discomfort or a burning sensation
Stop use and ask a doctor if you have
■ Rectal bleeding
■ No bowel movement after using this product
These may indicate a serious condition.
Keep out of reach of children
If swallowed, get medical help or contact a Poison Control Center right away.

Directions

■ Tear off one suppository from the strip
■ If the suppository seems soft, place it in the refrigerator before use or run cold water over it before removing the foil wrapper
■ Gently remove suppository by peeling apart the wrapper (see illustration 1)
■ Rub a water-based lubricant over the tip or dip it in water. It will help you slide it in smoothly
■ Lie on your side and insert 1 suppository, tapered end first, about the length of your finger into the rectum (see illustration 2)
■ Hold the suppository for about 15 - 20 minutes to allow time for it to melt
Adults and children 6 years of age and over 1 suppository once daily
Children under 6 years do not use

Other information

Keep away from excessive heat.

Inactive ingredients

purified water, sodium stearate.

Questions or Comments: call 1-844-474-7464


1

2




Front Foil (57237-339-51)

Annotation: Lot No. and Exp. Date will be embossed on each cavity on back foil as follows:
L.XXXXXXX
E.YYYY/MM



Back Foil (57237-339-51)



PROCTOZONE GMAX			
glycerin suppository			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57237-339
Route of Administration	RECTAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)		GLYCERIN	2 g
Inactive Ingredients			
Ingredient Name			Strength
WATER (UNII: 059QF0KO0R)			
SODIUM STEARATE (UNII: QU7E2XA9TG)			
Product Characteristics			
Color	white (clear)	Score	

Shape	BULLET		Size	
Flavor			Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57237-339-21	2 in 1 CARTON	07/31/2025	
1	NDC:57237-339-16	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:57237-339-52	5 in 1 CARTON	07/31/2025	
2	NDC:57237-339-51	5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M007.54	07/31/2025	

Labeler - Rising Pharma Holdings, Inc. (116880195)

Establishment

Name	Address	ID/FEI	Business Operations
Bliss GVS Pharma Limited		675630493	manufacture(57237-339) , analysis(57237-339) , pack(57237-339)

Revised: 7/2025

Rising Pharma Holdings, Inc.