

**DAYTIME SEVERE COLD AND FLU RELIEF SOFTGELS- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl capsule, liquid filled**  
**TopCo Associates LLC**

-----  
**Daytime Severe Cold and Flu Relief**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- ■ temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion & pressure ■ cough due to minor throat & bronchial irritation ■ minor aches & pains ■ headache ■ fever ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product,**

do not use more than directed.

**Stop use and ask a doctor if**

■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 softgels per 24 hours

---

adults & children 12 years & over	2 softgels with water every 4 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

---

- when using other Nighttime or Daytime products,  
carefully read each label to ensure correct dosing

**Other information**

- ■ store at room temperature

**Inactive ingredients**

FD&C Yellow # 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

**Questions or comments? 1-888-333-9792**

**PRINCIPAL DISPLAY PANEL**

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu\*

**MAXIMUM STRENGTH**

**Daytime Softgels**

**Severe Cold/Flu Relief**

ACETAMINOPHEN - Pain reliever; fever reducer

DEXTROMETHORPHAN HBr - Cough Suppressant

GUAIFENESIN- Expectorant

# PHENYLEPHRINE HCl - Nasal Decongestant

Relieves:

Headache, Fever, Sore throat, Minor aches & pains, Cough, Chest congestion, Nasal/Sinus congestion & Sinus pressure

Non-drowsy

Alcohol free

Antihistamine free

24 SOFTGELS

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER-EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

\*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademarks Vicks® DayQuil® Severe Cold & Flu.

24 SOFTGELS

TopCare health

MAXIMUM STRENGTH DayTime Cold & Flu Relief

SEVERE Cold & Flu Relief

**TopCare health**

MAXIMUM STRENGTH

## DayTime

### SEVERE Cold & Flu Relief

PAIN RELIEVER-FEVER REDUCER - ACETAMINOPHEN  
COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr  
EXPECTORANT - GUAIIFENESIN  
NASAL DECONGESTANT - PHENYLEPHRINE HCl

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Nasal/Sinus Congestion & Sinus Pressure
- Cough • Chest Congestion

Alcohol & Antihistamine Free  
Non-Drowsy

24 SOFTGELS actual size

COMBINE TO VICKS® DAYQUIL® SEVERE COLD & FLU ACTIVE INGREDIENTS\*

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER-EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

DISTRIBUTED BY  
TOPCO ASSOCIATES LLC  
ELK GROVE VILLAGE, IL 60007  
© TOPCO SPL1019  
QUESTIONS? 1-888-423-0139  
topcare@topco.com  
www.topcarebrand.com  
MADE IN INDIA

\*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademarks Vicks® DayQuil® Severe Cold & Flu.

Active ingredients (in each softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaiifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Drug Facts (continued)							
<p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves common cold/flu symptoms:                             <ul style="list-style-type: none"> <li>nasal congestion</li> <li>sinus congestion &amp; pressure</li> <li>cough due to minor throat &amp; bronchial irritation</li> <li>minor aches &amp; pains</li> <li>headache</li> <li>fever</li> <li>sore throat</li> </ul> </li> <li>reduces swelling of nasal passages</li> <li>temporarily restores freer breathing through the nose</li> <li>promotes nasal and/or sinus drainage</li> <li>helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 8 softgels in 24 hrs, which is the maximum daily amount for this product with other drugs containing acetaminophen. 3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>liver disease</li> <li>heart disease</li> <li>high blood pressure</li> <li>thyroid disease</li> <li>diabetes</li> <li>trouble</li> </ul>	<p>urinating due to enlarged prostate gland</p> <ul style="list-style-type: none"> <li>cough that occurs with too much phlegm (mucus)</li> <li>persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</b></p> <p><b>When using this product, do not use more than directed.</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>you get nervous, dizzy or sleepless</li> <li>pain, nasal congestion, or cough gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> </ul> <p><b>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning:</b> Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <p>take only as directed - see Overdose warning or do not exceed 8 softgels per 24 hours</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td>adults &amp; children 12 years &amp; over</td> <td>2 softgels with water every 4 hours</td> </tr> <tr> <td>children 4 to under 12 years</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 years</td> <td>do not use</td> </tr> </table> <p><b>When using other Nighttime or Daytime products, carefully read each label to ensure correct dosing.</b></p> <p><b>Other information</b> store at room temperature</p> <p><b>Inactive ingredients</b> FD&amp;C Red #40, FD&amp;C Yellow # 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p><b>Questions or comments?</b> 1-888-333-9792</p>	adults & children 12 years & over	2 softgels with water every 4 hours	children 4 to under 12 years	ask a doctor	children under 4 years	do not use
adults & children 12 years & over	2 softgels with water every 4 hours						
children 4 to under 12 years	ask a doctor						
children under 4 years	do not use						

Scan here for more information

LOT NO. \_\_\_\_\_ EXP. DATE: \_\_\_\_\_

# DAYTIME SEVERE COLD AND FLU RELIEF SOFTGELS

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-655
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	orange (Light Orange)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (oblong shaped)	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	341
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-655-24	2 in 1 CARTON	10/31/2019	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/31/2019	

**Labeler** - TopCo Associates LLC (006935977)

Revised: 12/2024

TopCo Associates LLC