STOOL SOFTENER LAXATIVE- docusate sodium and sennosides tablet P & L Development, LLC

Drug Facts

Active ingredients (in each tablet)

Docusate sodium 50 mg Sennosides 8.6 mg

Purpose

Stool softener

Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

- laxative products for longer than 1 week unless told to do so by a doctor
- if you are presently taking mineral oil, unless told to do so by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over a period of 2 weeks

Stop use and ask a doctor if

you have rectal bleeding or fail to have a bowel movement after use of a laxative.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• Take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses.

adults and children 12 years and older	take 2-4 tablets daily
children 6 to under 12 years of age	take 1-2 tablets daily
children 2 to under 6 years of age	take up to 1 tablet daily
children under 2	ask a doctor

Other information

- each tablet contains: calcium 30 mg
- each tablet contains: sodium 6 mg VERY LOW SODIUM
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, silcon dioxide, talc, titanium dioxide

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Colace® 2-IN-1*

Stool Softener + Stimulant Laxative

docusate sodium 50 mg

stool softener

sennosides 8.6 mg

stimulant laxative

- gentle, dependable
- relief in 6 to 12 hours

tablets


*This product is not manufactured of distributed by Avrio Health L.P., distributor of Colace® 2-IN1

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Distributed by: PL Developments

200 Hicks Street, Westbury, NY 11590

Product Label



Inactive ingredients croscarmellose sodiun dibasic calcium phosphate dihydrate, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, silicon dioxide, talc, tit dioxide Questions or comments?	Other information each tablet contains: c each tablet contains: s store at 25°C (77°F); ex 15-30°C (59-86°F)	children under 2	children 2 to under 6 years of age	children 6 to under 12 years of age	adults and children 12 years and over	Directions ■ take only by mouth. Dos daily dose, preferably in	hefore use. Keep out of reach of chill medical help or contact a least to the contact a le	Stop use and ask a doctor if you have or fail to have a bowel movement after. These may indicate a serious condition.	Ask a doctor before use if you have ■ stom ■ nausea ■ womiting ■ noticed a sudden c bowel habits that lasts over a period of 2 weeks	Drug Facts (continued)
S croscarmellose sodium, dihydrate, FD&C blue #2 #40 aluminum lake, stearate, microcrystalline col, silicon dioxide, talc, titanium ments?	calc sod	ask a doctor	take up to 1 tablet daily	take 1-2 tablets daily	take 2-4 tablets daily	lirections take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses.	medical help or contact a Polson Control Center right away.	Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.	if you have ■ stomach pain ■ noticed a sudden change in er a period of 2 weeks	red)

DOCUSATE SODIUM

50 mg

READYinCASE Stool Softener Laxative

STOOL SOFTENER LAXATIVE docusate sodium and sennosides tablet Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:59726-755 Route of Administration ORAL Active Ingredient/Active Moiety Ingredient Name Basis of Strength Strength

DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	

Product Characteristics					
Color	red	Score	no score		
Shape	ROUND	Size	10mm		
Flavor		Imprint Code	PSD21		
Contains					

l	Packaging							
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
	1	NDC:59726- 755-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/26/2021				

Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC Monograph Drug	505G(a)(3)	03/26/2021				

Labeler - P & L Development, LLC (800014821)

Revised: 9/2024 P & L Development, LLC