

ANTACID ANTIGAS MAXIMUM STRENGTH- aluminum hydroxide, magnesium hydroxide, simethicone liquid

Walgreens

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel USP) 800 mg

Magnesium Hydroxide 800 mg

Simethicone 80 mg

Purpose

Antacid

Antacid

Antigas

Uses

- for the relief of
 - acid indigestion
 - heartburn
 - sour stomach
 - upset stomach associated with these symptoms
 - pressure and bloating commonly referred to as gas
 - overindulgence in food and drink

Warnings

Do not take

more than 60 mL in a 24-hour period or use the maximum dosage for more than 2 weeks except under the advice and supervision of a physician

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

symptoms last more than 2 weeks.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- shake well before using
- mL = milliliter
- do not take more than 60 mL in 24 hours or use the maximum dosage for more than 2 weeks
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and older: take 10-20 mL as needed, between or after meals, at bedtime, or as directed by a doctor
- children under 12 years of age: do not use

Other information

- **each 10 mL contains:** magnesium 335 mg, sodium 10 mg
- store between 20-25°C (68-77°F). Avoid freezing

Inactive ingredients

benzyl alcohol, carboxymethylcellulose sodium, flavor, glycerin, microcrystalline cellulose, purified water, sodium carbonate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Mylanta® Maximum Strength Active Ingredients^{††}

Antacid +

Anti-gas Liquid

ALUMINUM HYDROXIDE 800 mg / ANTACID

MAGNESIUM HYDROXIDE 800 mg / ANTACID

SIMETHICONE 80 mg / ANTI-GAS

MAXIMUM STRENGTH

- Fast relief of heartburn, pressure, bloating, acid indigestion & sour stomach

CLASSIS FLAVOR

^{††}This product is not manufactured or distributed by McNeil Consumer Pharmaceuticals Co., distributor of Mylanta® Maximum Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO.,

200 WILMOT RD., DEERFIELD, IL 60015

walgreens.com

Walgreens

Compare to Mylanta® Maximum Strength Active Ingredients††

NDC 0363-0528-12

Antacid + Anti-Gas Liquid

ALUMINUM HYDROXIDE 800 mg / ANTACID
MAGNESIUM HYDROXIDE 800 mg / ANTACID
SIMETHICONE 80 mg / ANTI-GAS

MAXIMUM STRENGTH

- Fast relief of heartburn, pressure, bloating, acid indigestion & sour stomach

12 FL OZ (355 mL)

CLASSIC FLAVOR

PLD-A4608 LB905706

Active ingredients (in each 10 mL)	Purposes
Aluminum hydroxide [equiv. to dried gel USP] 800 mg	Antacid
Magnesium hydroxide 800 mg	Antacid
Simethicone 80 mg	Antigas

Uses ■ for the relief of

- acid indigestion
- heartburn
- sour stomach
- upset stomach associated with these symptoms
- pressure and bloating commonly referred to as gas
- overindulgence in food and drink

Do not take more than 60 mL in a 24-hour period or use the maximum dosage for more than 2 weeks except under the advice and supervision of a physician.

- kidney disease
- a magnesium-restricted diet

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

¹¹ This product is not manufactured or distributed by McNeil Consumer Pharmaceuticals Co., distributor of Mylanta® Maximum Strength.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2018 Walgreen Co.

ITEM 272294 W10023-0418-F



ORG0418-F

PLD-A460B LB005707

PEEL CORNER FOR MORE DRUG FACTS

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

- shake well before using
- mL = milliliter
- do not take more than 60 mL in 24 hours or use the maximum dosage for more than 2 weeks
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years of age and older: take 10-20 mL as needed, between or after meals, at bedtime, or as directed by a doctor
- children under 12 years of age: do not use

- each 10 mL contains: magnesium 335 mg, sodium 10 mg
- store between 20-25°C (68-77°F). Avoid freezing.

carboxymethylcellulose sodium, flavor, glycerin, microcrystalline cellulose, purified water, sodium carbonate, sorbitol, sucralose, xanthan gum

Call 1-877-753-3935 Monday-Friday
9AM-5PM EST

aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0528
Route of Administration	ORAL		

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	800 mg in 10 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients

Ingredient Name				Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)				
GLYCERIN (UNII: PDC6A3C0OX)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
WATER (UNII: 059QF0KO0R)				
SODIUM CARBONATE (UNII: 45P3261C7T)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0528-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL		part331	03/31/2018	

Labeler - Walgreens (008965063)

Revised: 10/2019

Walgreens