


**DIPYRIDAMOLE- dipyridamole tablet**  
**Amneal Pharmaceuticals of New York LLC**

-----  
**Dipyridamole Tablets, USP**  
**(25 mg, 50 mg and 75 mg)**  
**Rx only**

**DESCRIPTION**

Dipyridamole is a platelet inhibitor chemically described as 2,2',2'',2'''-[(4,8-Dipiperidinopyrimido[5,4-*d*]pyrimidine-2,6-diyl)dinitrilo]-tetraethanol. The molecular weight is 504.63 and the molecular formula is C<sub>24</sub>H<sub>40</sub>N<sub>8</sub>O<sub>4</sub>. The structural formula is represented below:



Dipyridamole, USP is intensely yellow crystalline powder or needles. It is practically insoluble in water, sparingly soluble in ethyl alcohol, very slightly soluble in acetone and ethyl acetate.

Each tablet, for oral administration, contains 25 mg, 50 mg or 75 mg dipyridamole, USP. In addition, each tablet contains the following inactive ingredients: colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, Type A, talc, and titanium dioxide.

**CLINICAL PHARMACOLOGY**

It is believed that platelet reactivity and interaction with prosthetic cardiac valve surfaces, resulting in abnormally shortened platelet survival time, is a significant factor in thromboembolic complications occurring in connection with prosthetic heart valve replacement.

Dipyridamole tablets have been found to lengthen abnormally shortened platelet survival time in a dose-dependent manner.

In three randomized controlled clinical trials involving 854 patients who had undergone surgical placement of a prosthetic heart valve, dipyridamole tablets, in combination with warfarin, decreased the incidence of postoperative thromboembolic events by 62% to

91% compared to warfarin treatment alone. The incidence of thromboembolic events in patients receiving the combination of dipyridamole tablets and warfarin ranged from 1.2% to 1.8%. In three additional studies involving 392 patients taking dipyridamole tablets and coumarin-like anticoagulants, the incidence of thromboembolic events ranged from 2.3% to 6.9%.

In these trials, the coumarin anticoagulant was begun between 24 hours and 4 days postoperatively, and the dipyridamole tablets were begun between 24 hours and 10 days postoperatively. The length of follow-up in these trials varied from 1 to 2 years.

Dipyridamole tablets do not influence prothrombin time or activity measurements when administered with warfarin.

### **Mechanism of Action**

Dipyridamole inhibits the uptake of adenosine into platelets, endothelial cells and erythrocytes *in vitro* and *in vivo*; the inhibition occurs in a dose-dependent manner at therapeutic concentrations (0.5 mcg/mL to 1.9 mcg/mL). This inhibition results in an increase in local concentrations of adenosine which acts on the platelet A<sub>2</sub>-receptor thereby stimulating platelet adenylate cyclase and increasing platelet cyclic-3',5'-adenosine monophosphate (cAMP) levels. Via this mechanism, platelet aggregation is inhibited in response to various stimuli such as platelet activating factor (PAF), collagen and adenosine diphosphate (ADP).

Dipyridamole inhibits phosphodiesterase (PDE) in various tissues. While the inhibition of cAMP-PDE is weak, therapeutic levels of dipyridamole inhibit cyclic-3',5'-guanosine monophosphate-PDE (cGMP-PDE), thereby augmenting the increase in cGMP produced by EDRF (endothelium-derived relaxing factor, now identified as nitric oxide).

### **Hemodynamics**

In dogs intraduodenal doses of dipyridamole of 0.5 mg/kg to 4.0 mg/kg produced dose-related decreases in systemic and coronary vascular resistance leading to decreases in systemic blood pressure and increases in coronary blood flow. Onset of action was in about 24 minutes and effects persisted for about 3 hours.

Similar effects were observed following intravenous dipyridamole in doses ranging from 0.025 mg/kg to 2.0 mg/kg.

In man the same qualitative hemodynamic effects have been observed. However, acute intravenous administration of dipyridamole may worsen regional myocardial perfusion distal to partial occlusion of coronary arteries.

### **Pharmacokinetics and Metabolism**

Following an oral dose of dipyridamole tablets, the average time to peak concentration is about 75 minutes. The decline in plasma concentration following a dose of dipyridamole tablets fits a two-compartment model. The alpha half-life (the initial decline following peak concentration) is approximately 40 minutes. The beta half-life (the terminal decline in plasma concentration) is approximately 10 hours. Dipyridamole is highly bound to plasma proteins. It is metabolized in the liver where it is conjugated as a glucuronide and excreted with the bile.

## **INDICATIONS AND USAGE**

Dipyridamole tablets are indicated as an adjunct to coumarin anticoagulants in the prevention of postoperative thromboembolic complications of cardiac valve replacement.

## **CONTRAINDICATIONS**

Hypersensitivity to dipyridamole and any of the other components.

## **PRECAUTIONS**

### **General**

*Coronary Artery Disease:* Dipyridamole has a vasodilatory effect and should be used with caution in patients with severe coronary artery disease (e.g., unstable angina or recently sustained myocardial infarction). Chest pain may be aggravated in patients with underlying coronary artery disease who are receiving dipyridamole.

*Hepatic Insufficiency:* Elevations of hepatic enzymes and hepatic failure have been reported in association with dipyridamole administration.

*Hypotension:* Dipyridamole should be used with caution in patients with hypotension since it can produce peripheral vasodilation.

*Stress Testing with Intravenous Dipyridamole and Other Adenosinergic Agents:* Clinical experience suggests that patients being treated with dipyridamole tablets who also require pharmacological stress testing with intravenous dipyridamole or other adenosinergic agents (e.g. adenosine, regadenoson) should interrupt dipyridamole tablets for 48 hours prior to stress testing.

Intake of dipyridamole tablets within 48 hours prior to stress testing with intravenous dipyridamole or other adenosinergic agents may increase the risk for cardiovascular side effects of these agents and may impair the sensitivity of the test.

### **Laboratory Tests**

Dipyridamole has been associated with elevated hepatic enzymes.

### **Drug Interactions**

No pharmacokinetic drug-drug interaction studies were conducted with dipyridamole tablets. The following information was obtained from the literature.

*Adenosinergic agents (e.g., adenosine, regadenoson):* Dipyridamole has been reported to increase the plasma levels and cardiovascular effects of adenosine. Adjustment of adenosine dosage may be necessary. Dipyridamole also increases the cardiovascular effects of regadenoson, an adenosine A<sub>2A</sub>-receptor agonist. The potential risk of cardiovascular side effects with intravenous adenosinergic agents may be increased during the testing period when dipyridamole is not held 48 hours prior to stress testing.

*Cholinesterase Inhibitors:* Dipyridamole may counteract the anticholinesterase effect of cholinesterase inhibitors, thereby potentially aggravating myasthenia gravis.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

In studies in which dipyridamole was administered in the feed to mice (up to 111 weeks in males and females) and rats (up to 128 weeks in males and up to 142 weeks in females), there was no evidence of drug-related carcinogenesis. The highest dose administered in these studies (75 mg/kg/day) was, on a mg/m<sup>2</sup> basis, about equivalent to the maximum recommended daily human oral dose (MRHD) in mice and about twice the MRHD in rats. Mutagenicity tests of dipyridamole with bacterial and mammalian cell systems were negative. There was no evidence of impaired fertility when dipyridamole was administered to male and female rats at oral doses up to 500 mg/kg/day (about 12 times the MRHD on a mg/m<sup>2</sup> basis). A significant reduction in number of corpora lutea with consequent reduction in implantations and live fetuses was, however, observed at 1,250 mg/kg (more than 30 times the MRHD on a mg/m<sup>2</sup> basis).

## **Pregnancy**

### ***Teratogenic Effects***

Reproduction studies have been performed in mice, rabbits and rats at oral dipyridamole doses of up to 125 mg/kg, 40 mg/kg and 1,000 mg/kg, respectively (about 1½, 2 and 25 times the maximum recommended daily human oral dose, respectively, on a mg/m<sup>2</sup> basis) and have revealed no evidence of harm to the fetus due to dipyridamole. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, dipyridamole tablets should be used during pregnancy only if clearly needed.

## **Nursing Mothers**

As dipyridamole is excreted in human milk, caution should be exercised when dipyridamole tablets are administered to a nursing woman.

## **Pediatric Use**

Safety and effectiveness in the pediatric population below the age of 12 years have not been established.

## **ADVERSE REACTIONS**

Adverse reactions at therapeutic doses are usually minimal and transient. On long-term use of dipyridamole tablets initial side effects usually disappear. The following reactions in Table 1 were reported in two heart valve replacement trials comparing dipyridamole tablets and warfarin therapy to either warfarin alone or warfarin and placebo:

**Table 1: Adverse Reactions Reported in 2 Heart Valve Replacement Trials**

<b>Adverse Reaction</b>	<b>Dipyridamole Tablets/ Warfarin</b>	<b>Placebo/ Warfarin</b>
Number of patients	147	170
Dizziness	13.6%	8.2%
Abdominal distress	6.1%	3.5%
Headache	2.3%	0.0%
Rash	2.3%	1.1%

Other reactions from uncontrolled studies include diarrhea, vomiting, flushing and pruritus. In addition, angina pectoris has been reported rarely and there have been rare reports of liver dysfunction. On those uncommon occasions when adverse reactions have been persistent or intolerable, they have ceased on withdrawal of the medication.

When dipyridamole tablets were administered concomitantly with warfarin, bleeding was no greater in frequency or severity than that observed when warfarin was administered alone. In rare cases, increased bleeding during or after surgery has been observed.

In post-marketing reporting experience, there have been rare reports of hypersensitivity reactions (such as rash, urticaria, severe bronchospasm, and angioedema), larynx edema, fatigue, malaise, myalgia, arthritis, nausea, dyspepsia, paresthesia, hepatitis, thrombocytopenia, alopecia, cholelithiasis, hypotension, palpitation, and tachycardia.

## **OVERDOSAGE**

In case of real or suspected overdose, seek medical attention or contact a Poison Control Center immediately. Careful medical management is essential. Based upon the known hemodynamic effects of dipyridamole, symptoms such as warm feeling, flushes, sweating, restlessness, feeling of weakness and dizziness may occur. A drop in blood pressure and tachycardia might also be observed.

Symptomatic treatment is recommended, possibly including a vasopressor drug. Gastric lavage should be considered. Administration of xanthine derivatives (e.g., aminophylline) may reverse the hemodynamic effects of dipyridamole overdose. Since dipyridamole is highly protein bound, dialysis is not likely to be of benefit.

## **DOSAGE AND ADMINISTRATION**

### *Adjunctive Use in Prophylaxis of Thromboembolism after Cardiac Valve Replacement.*

The recommended dose is 75 mg to 100 mg four times daily as an adjunct to the usual warfarin therapy. Please note that aspirin is not to be administered concomitantly with coumarin anticoagulants.

## **HOW SUPPLIED**

Dipyridamole Tablets USP, **25 mg** are white to pale yellow, round, standard convex film-coated tablets debossed with "C81" on one side and plain on the other side.

They are available as follows:

Bottles of 100: NDC 0115-1070-01

Bottles of 1,000: NDC 0115-1070-03

Dipyridamole Tablets USP, **50 mg** are white to pale yellow, round, standard convex film-coated tablets debossed with "C82" on one side and plain on the other side.

They are available as follows:

Bottles of 100: NDC 0115-1071-01

Bottles of 1,000: NDC 0115-1071-03

Dipyridamole Tablets USP, **75 mg** are white to pale yellow, round, standard convex film-coated tablets debossed with "C83" on one side and plain on the other side.

They are available as follows:

Bottles of 100: NDC 0115-1072-01

Bottles of 1,000: NDC 0115-1072-03

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Keep this and all medication out of the reach of children.

Dispense in tightly-closed, light-resistant container as defined in the USP, with child-resistant closure, as required.

**To report SUSPECTED ADVERSE REACTIONS, contact Amneal Pharmaceuticals at 1-877-835-5472 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

Manufactured by:  
**Amneal Pharmaceuticals Pvt. Ltd.**  
**Oral Solid Dosage Unit**  
Ahmedabad 382213, INDIA

Distributed by:  
**Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807

Rev. 08-2021-00

### PRINCIPAL DISPLAY PANEL - 25 mg Tablet Label

The image shows a rectangular label for Dipyridamole Tablets, USP, 25 mg. The label is divided into several sections. On the left, there is a purple banner with the text "25 mg" in white. Below this, the Amneal logo is visible. The top left corner contains the NDC number "NDC 0115-1070-01". The main text reads "Dipyridamole Tablets, USP" in a large, bold font. Below this, it says "Rx only" and "100 Tablets". The right side of the label contains detailed information: "Each tablet contains: Dipyridamole, USP .....25 mg", "USUAL DOSAGE: See accompanying outsert for additional dosing information.", "Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].", "Do not use if printed safety seal under cap is broken or missing.", "Keep this and all drugs out of the reach of children.", "Manufactured by: Amneal Pharmaceuticals Pvt. Ltd. Oral Solid Dosage Unit Ahmedabad 382213, INDIA", "Distributed by: Amneal Pharmaceuticals LLC Bridgewater, NJ 08807", "Rev. 08-2021-00", and "Mfg. Lic. No. G/25/2137". A barcode is located on the right side of the label, with the number "3 01151 07001 1" printed below it. On the far right, there is a vertical text area that reads "Non-Varnish Area (For Lot And Exp. Date) (26 X 28 mm)".


### PRINCIPAL DISPLAY PANEL - 50 mg Tablet Label

NDC 0115-1071-01

# Dipyridamole Tablets, USP

**50 mg**

Rx only  
100 Tablets



Each tablet contains: Dipyridamole, USP .....50 mg  
**USUAL DOSAGE:** See accompanying outsert for additional dosing information.  
**Store at 20° to 25°C (68° to 77°F)** [see USP Controlled Room Temperature].  
**Do not use if printed safety seal under cap is broken or missing.**  
 Keep this and all drugs out of the reach of children.  
 Manufactured by: **Amneal Pharmaceuticals Pvt. Ltd.**  
**Oral Solid Dosage Unit**  
 Ahmedabad 382213, INDIA  
 Distributed by: **Amneal Pharmaceuticals LLC**  
 Bridgewater, NJ 08807  
 Rev. 08-2021-00 Mfg. Lic. No. G/25/2137



3 01151 07101 8

Non-Varnish Area  
(For Lot And Exp. Date)  
(26 X 28 mm)


## PRINCIPAL DISPLAY PANEL - 75 mg Tablet Label

NDC 0115-1072-01


# Dipyridamole Tablets, USP

**75 mg**

Rx only  
100 Tablets



Each tablet contains: Dipyridamole, USP .....75 mg  
**USUAL DOSAGE:** See accompanying outsert for additional dosing information.  
**Store at 20° to 25°C (68° to 77°F)** [see USP Controlled Room Temperature].  
**Do not use if printed safety seal under cap is broken or missing.**  
 Keep this and all drugs out of the reach of children.  
 Manufactured by: **Amneal Pharmaceuticals Pvt. Ltd.**  
**Oral Solid Dosage Unit**  
 Ahmedabad 382213, INDIA  
 Distributed by: **Amneal Pharmaceuticals LLC**  
 Bridgewater, NJ 08807  
 Rev. 08-2021-00 Mfg. Lic. No. G/25/2137



3 01151 07201 5

Non-Varnish Area  
(For Lot And Exp. Date)  
(26 X 28 mm)

## DIPYRIDAMOLE

dipyridamole tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0115-1070
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIPYRIDAMOLE</b> (UNII: 64ALC7F90C) (DIPYRIDAMOLE - UNII:64ALC7F90C)	DIPYRIDAMOLE	25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	white (white to pale yellow)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	C81
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0115-1070-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/30/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040782	08/30/2021	

## DIPYRIDAMOLE

dipyridamole tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0115-1071
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIPYRIDAMOLE</b> (UNII: 64ALC7F90C) (DIPYRIDAMOLE - UNII:64ALC7F90C)	DIPYRIDAMOLE	50 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	



<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	white (white to pale yellow)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	C82
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0115-1071-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/30/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040782	08/30/2021	

## DIPYRIDAMOLE

dipyridamole tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0115-1072
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIPYRIDAMOLE</b> (UNII: 64ALC7F90C) (DIPYRIDAMOLE - UNII:64ALC7F90C)	DIPYRIDAMOLE	75 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	white (white to pale yellow)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	C83
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0115-1072-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/30/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040782	08/30/2021	

**Labeler** - Amneal Pharmaceuticals of New York LLC (123797875)

### Establishment

Name	Address	ID/FEI	Business Operations
Amneal Pharmaceuticals Private Limited		650762060	analysis(0115-1070, 0115-1071, 0115-1072) , label(0115-1070, 0115-1071, 0115-1072) , manufacture(0115-1070, 0115-1071, 0115-1072) , pack(0115-1070, 0115-1071, 0115-1072)

Revised: 4/2023

Amneal Pharmaceuticals of New York LLC