## NICOTINE POLACRILEX- nicotine polacrilex gum, chewing Lucy Goods, Inc.

-----

Nicotine Polacrilex Gum 2 mg and 4 mg Coated Fruit - Rugby

#### Drug Facts

#### Active ingredient (in each chewing piece)

Nicotine Polacrilex 2 mg (nicotine)

#### Purpose

Stop smoking aid

#### Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

#### Warnings

**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

#### Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

#### Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

#### Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

#### Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions - 2 mg

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every	1 piece every	1 piece every
1 to 2 hours	2 to 4 hours	4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

#### Other information - 2 mg

- each piece contains: calcium 115 mg, sodium 8 mg
- store at 20 25ºC (68 77ºF)
- protect from light

#### Inactive ingredients - 2 mg

acacia, acesulfame potassium, carnauba wax, FD&C blue #2 lake, FD&C red #40, FD&C yellow #6 lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide, xylitol.

#### **Questions or comments?**

Call

#### Principal display panel - 2 mg

\*Compare to the active ingredient of Nicorette® Gum

Nicotine Gum

Nicotine polacrilex gum USP, 2 mg (nicotine)

Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE **MORE THAN 30 MINUTES** AFTER WAKING UP

If you smoke your first cigarette <u>WITHIN</u> 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

2 mg

Sugar-Free

COATED FRUIT

#### TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivated to quit.
- 2. Use Enough Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
- 3. Use Long Enough Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. Use with a Support Program as directed in the enclosed User's Guide.
- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

## BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

\*This product is not manufactured or distributed by Haleon Group of Companies, distributor of Nicorette® Gum.

#### Marketed By: Lucy Goods Inc.

2510 E Sunset Road

Suite 5 #A6546

Las Vegas, NV 89120


### Package label 2 mg

#### TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.

- Use Enough Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
  Use Long Enough Use Nicotine Polacrilex Gum for the full 12 weeks.
- Use With a Support Program as directed in the enclosed User's Guide.







\*Compare to the active ingredient in Nicorette\* Gum

FRUIT

Actual Size

# **Nicotine Gum**

Nicotine polacrilex gum USP, 2 mg (nicotine) Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP

If you smoke your first cigarette <u>WITHIN</u> 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg



Sugar-Free

I

100 Pieces, 2 mg EACH



# Nicotine Gum

Nicotine polacrilex gum USP, 2 mg (nicotine) Stop Smoking Aid

6 72552 69005 8	
2510 E Sunaet Road Suite 5 #A656 Las Vegas, VV ,segeV at	Directions at you arenable years of age, and a decire before the comparison a boyn using this product, react the enclosed bears Daule the comparison a boyn using this year. This digwethe within 30 minutes of waiding up, use at year and they trayer the figure of they at year and they are the second they at year and they are the second at year and the second the second at year and the second the second at year and the second the second the second at year and the second the second the second the second at year and the second the attemption of the second
"The product is not insurface to distributed by Hellow group of comparises. distribute of Noosate" Gun. Mar Keet ed. By : Lucy Goods Inc.	Kong yan i na kata kata kata ya kata y Kata ya kata ya kata ya kata ya kata ya kata ya kata ya kata
residences in more could's of bet-direct to be due to a propose of T	you have symptoms of an allergic reaction (such as difficulty breathing or reat).



00 Pieces, 2 mg EACH	EACH	6w 7	Pieces,	001
----------------------	------	------	---------	-----



1TON-1



109120	Stop use and ask a doctor il mouth, tooft, or juw proleme occu a imegular hourboot or pajolitatore cocur	FOSC DEB 12 MM, FOSC	og arrethæres, accester <b>21</b> 184 welley 33671,046 ber 1 multos , stelan mulsarger	, Seed mug , Shoreft , Evis
If you among your first cigan the second and se second and second and seco	have a doctave or plasmaactur before used if you am a taakay presentificion madicare for depression or satimma. Your prescription does a taakay or one doub satis and doub and taakay or one doub satis and taakay or one of taakay and taakay or one doub satis and taakay or one of taakay and taakay or one doub satis and taakay or one of taakay and taakay or one of taakay or one of taakay or one of taakay and taakay or one of taakay or one of taakay or one of taakay or one of taakay and taakay or one of taakay and taakay or one of taakay or one of taakay or one of taakay or one of taakay or one of t	Other information store st 20 - 25°C (68 - 7 store st 20 - 25°C (68 - 7 store st 20 - 15°C (68 - 7 store st 700 light	8 multipos gan 211 multiplas	<u>fu</u>
(entrovin) gm S, 9SU mug valiccelog entrovi Stop Smoking bid Foa Troose wwo swoke Thera Plast crowner Te Plast Wown ST No 25 No 1990 W	e with use of the background of the back of the background of the background of the background of the back determined of the background of	May a manufacture strang within the hour Howew within the hour Howew since this may cause y and ont use more than 2 and and use more than a the important to complete a	6 pieces a day or frequent craxings, you no frequent breatburn, neu- treatburn, neu- treatburn, neu- treatburn, you or frequent breatburn, neu- treatburn, neu-	n nead is use it second piece are proce after arceler ea or other side aftects.
Micotine Gum	Asympton Believed to be earlier an provides. Stavids provide the medicate on the advice ad your feedbar and or to the stavid provide the advice of the and of the advice ad your feedbar and and a stavid provide the advice of the and advice a advice and and the advice advi	These is the second of the sec		
	<b>1680</b> nitro benational generation protocol protocol and the security devices protocol and the protocol of the	t to 2 terms and the second of	2 to 4 hours the used a co 2 to 4 hours 2 hours the used a co	utariu waak po diat pue poet 4 po 8 pontus 1 biece sweuk
	bie grokerna gota	8 of 1 askell	6 of 7 syseew	Weeks 10 to 12
TUCK	esognu <sup>q</sup> (acab priverio doea ni) traibangni avitos		DE matt enom ethenegio i privedal ett of gritneaae	,qu gobiew nothe sofuniar shidorice Xeew ST
	spey brug	Drug Facts (contin	(par	100m Oct. 544

LED-VEILER FC006333

NEOL NO AGEO GUY GTAGS TATOMONI Al Jac Don not toolitali manga mga dogayagaa

#### **LUCY Nicotine Gum Coated Fruit**

nicotine polacrilex gum, chew	ing				
Product Information					
Product Type	HUMAN OTC DRUG	Item Code	(Source)	NDC:72	2224-925
Route of Administration	ORAL				
Active Ingredient/Active	Moiety				
-	ient Name		Basis of Stre	ngth	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NI		k)	NICOTINE		2 mg
Inactive Ingredients					
mactive myreulents	Ingredient Name				Strength
ACACIA (UNII: 5C5403N260)	ingredient Name				Strength
ACESULFAME POTASSIUM (UNII:	230\/730569)				
CARNAUBA WAX (UNII: R12CBM0E					
FUQL BLUE NU. 2 (UNII: LUDKOK/					
FD&C BLUE NO. 2 (UNII: L06K8R7 FD&C RED NO. 40 (UNII: WZ B912					
FD&C RED NO. 2 (UNII: LUGKAR7 FD&C RED NO. 40 (UNII: WZ B912 FD&C YELLOW NO. 6 (UNII: H77V	7XOA)				
FD&C RED NO. 40 (UNII: WZ B912	7XOA) EI93A8)	16N6OH)			
FD&C RED NO. 40 (UNII: WZB912 FD&C YELLOW NO. 6 (UNII: H77V	7XOA) EI93A8) JNSPECIFIED (UNII: 9XZ8H	16N6OH)			
FD&C RED NO. 40 (UNII: WZB912 FD&C YELLOW NO. 6 (UNII: H77V HYDROXYPROPYL CELLULOSE, U MAGNESIUM OXIDE (UNII: 3A3U00	7XOA) EI93A8) J <b>NSPECIFIED</b> (UNII: 9XZ8F GI71G)	16N6OH)			
FD&C RED NO. 40 (UNII: WZB912 FD&C YELLOW NO. 6 (UNII: H77V HYDROXYPROPYL CELLULOSE, U MAGNESIUM OXIDE (UNII: 3A3U00 SODIUM BICARBONATE (UNII: 8M	7XOA) EI93A8) JNSPECIFIED (UNII: 9XZ8H GI71G) DF5V39QO)	16N6OH)			
FD&C RED NO. 40 (UNII: WZ B912 FD&C YELLOW NO. 6 (UNII: H77V HYDROXYPROPYL CELLULOSE, U MAGNESIUM OXIDE (UNII: 3A3U00 SODIUM BICARBONATE (UNII: 8M SODIUM CARBONATE (UNII: 45P3	7XOA) EI93A8) JNSPECIFIED (UNII: 9XZ8H GI71G) DF5V39QO)	16N6OH)			
FD&C RED NO. 40 (UNII: WZ B912) FD&C YELLOW NO. 6 (UNII: H77V) HYDROXYPROPYL CELLULOSE, U MAGNESIUM OXIDE (UNII: 3A3U00) SODIUM BICARBONATE (UNII: 8M) SODIUM CARBONATE (UNII: 45P3) SUCRALOSE (UNII: 96K6UQ3Z D4)	7XOA) EI93A8) JNSPECIFIED (UNII: 9XZ8H GI71G) DF5V39QO)	16N6OH)			
FD&C RED NO. 40 (UNII: WZB912 FD&C YELLOW NO. 6 (UNII: H77V HYDROXYPROPYL CELLULOSE, U	7XOA) EI93A8) JNSPECIFIED (UNII: 9XZ8F GI71G) DF5V39QO) 261C7T)	16N6OH)			

Pr	oduct Chara	cteristics		
Co	lor	brown (Light Brown)	core	no score
Sh	аре	SQUARE S	Size	14mm
Fla	avor	FRUIT	mprint Code	
Со	ntains			
Pa	ackaging			
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:72224-925- 10	100 in 1 CARTON	04/30/2025	
1		1 in 1 BLISTER PACK; Type 0: Not a Combinatio Product	n	
	arketing I	nformation		
Μ			n Marketing Start	Marketing End
M	Marketing Category	Application Number or Monograpl Citation	Date	Date

## Labeler - Lucy Goods, Inc. (080875415)

Revised: 4/2025

Lucy Goods, Inc.