#### SENSICAINE TM EXTRA TOPICAL ANESTHETIC- benzocaine gel Safco Dental Supply Co.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### SensiCaine<sup>TM</sup> Extra Topical Anesthetic Gel Mint Flavor

#### Active ingridients (in each gram)

Benzocaine 200 mg

#### Purpose

Oral Anesthetic

#### Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

#### Warnings

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** Do not use on patients with a history of allergies to local anesthetics such as benzocaine or other "caine" anesthetics.

**Sore throat warning:**If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsen, see your dentist or doctor promptly.

#### Do not use

- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

If pregnant or breast feeding, consult a doctor before use.

**KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center immediately.

#### Directions

- adults and children 2 years of age and older: apply only amount needed to the oral mucosa to prevent or relieve pain
- children under 2 years of age: do not use

#### Other information

- Store at 59 86°F (15 30°C).
- Protect from freezing.

#### **Inactive ingredients**

flavoring, PEG 3350, PEG 400, sucralose. May contain D&C Red #33, FD&C Red #40, D&C Green #5, FD&C Yellow #5 as a color additive.

#### Questions or comments?

800-621-2178 www.safcodental.com

#### **PRINCIPAL DISPLAY PANEL - Jar Label**

SAFCO SensiCaine<sup>TM</sup> Extra

Topical Anesthetic Gel

Contains 20% Benzocaine

#### **Mint Flavor**

For Professional Use Only

1 oz. (34 g)



## **Drug Facts**

### Active ingredients (in each gram)

Benzocaine 200 mg

# Purpose

Oral Anesthetic

### Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

# Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: pale, gray, or blue colored skin (cyanosis) headache rapid heart rate shortness of breath dizziness or lightheadedness

•fatigue or lack of energy

Allergy alert: Do not use on patients with a history of allergies to local anesthetics such as benzocaine or other fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsen, see your dentist or doctor promptly.

#### Do not use •for teething

•in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

If pregnant or breast feeding, consult a doctor before use.

KEEP OUT OF REACH OF CHILDREN. In case of overdose, get medical help or contact a Poison Control Center immediately.

#### **Directions** •adults and children 2 years of age and older: apply only amount needed to the oral mucosa to prevent or relieve pain •children under 2 years of age: do not use

**Other information** •Store at 59 - 86°F (15 - 30°C). •Protect from freezing.

**Inactive ingredients** flavoring, PEG 3350, PEG 400, sucralose. May contain D&C Red #33, FD&C Red #40, D&C Green #5, FD&C Yellow #5 as a color additive.

Questions or

"caine" anesthetics. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by



#### SENSICAINE TM EXTRA TOPICAL ANESTHETIC

benzocaine gel

Product Information									
Product Type		HUMAN OTC DRUG	Ite	m Code (S	Code (Source)		NDC:67239-0213		
Route of Administrati	ion	DENTAL							
Active Ingredient/Active Moiety									
	Ingre	dient Name		Basis of Strengtl		Strength			
Benzocaine (UNII: U3RS	SY48JW5) (Ben	zocaine - UNII:U3RSY48JW5)			Benzocaine		200 mg in 1 g		
<b>Inactive Ingredien</b>	nts								
Ingredient Name							Strength		
Polyethylene Glycol 3350 (UNII: G2M7P15E5P)									
Polyethylene Glycol 400 (UNII: B697894SGQ)									
Sucralose (UNII: 96K6UQ3ZD4)									
D&C RED NO.33 (UNII: 9DBA0SBB0L)									
FD&C RED NO.40 (UNII: WZB9127XOA)									
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)									
FD&C YELLOW NO.5 (UNII: I753WB2F1M)									
Product Character	ristics								
Color		green	Score						
Shape			Size						
Flavor		MINT	Imprint Code						
Contains									
Packaging									
# Item Code	Package Description				ting Start Date	Mark	eting End Date		
1 NDC:67239-0213-1	29 g in 1 JAR; Type 0: Not a Combination Product 04/30			04/30/20	17				

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC MONOGRAPH NOT FINAL	part356	04/30/2017					
[	<b>F</b>						

Labeler - Safco Dental Supply Co. (020417380)

Registrant - Safco Dental Supply Co. (020417380)

Revised: 11/2018

Safco Dental Supply Co.