

## **BASIC CARE MIGRAINE RELIEF- acetaminophen, aspirin, caffeine tablet, film coated**

**Amazon.com Services LLC**

-----  
**Amazon Migraine Relief Drug Facts**

### **Active ingredients (in each tablet)**

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

### **Purposes**

Pain reliever

Pain reliever aid

### **Use**

- treats migraine

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

you take

- more than 2 gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause severe nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever

reducers

- you have vomiting with migraine headache

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
- diabetes
- gout
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

### **stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not use more than directed**
- adults: take 2 geltabs with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 geltabs in 24 hours, unless directed by a doctor
- under 18 years of age: as a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

### **Inactive ingredients**

black iron oxide, carnauba wax, croscopovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, gelatin, glycerin, hypromellose, maltitol solution, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polyvinyl acetate phthalate, povidone, pregelatinized starch, propylene glycol, stearic acid, titanium dioxide

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

amazon

basic care

Treats Migraine

Compare to Excedrin<sup>®</sup> Migraine active ingredients

Migraine Relief

Acetaminophen/Aspirin (NSAID)/Caffeine Tablets

Pain Reliever/Pain Reliever Aid

actual size

80 GELTABS\*\*

\*\*Gelatin Coated Tablets

22527 BN C1



**Drug Facts**

**Active ingredients**  
(In each geltab)  
Acetaminophen 250 mg, Aspirin 250 mg (NSAID), Caffeine 65 mg

**Pain reliever**  
Pain reliever aid  
non-steroidal anti-inflammatory drug

**Use** ■ treats migraine

**Warnings**

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin redness ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
Allergy alert: A skin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ stomach pain ■ difficulty breathing ■ dizziness ■ difficulty swallowing  
Liver damage may occur if you take  
■ more than 2 tablets in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
Some of the following warning signs may cause severe stomach bleeding. The chance is higher if you  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ have had stomach or intestinal surgery or stomach drug  
■ take other drugs containing aspirin or nonprescription NSAIDs (ibuprofen, naproxen, or others)  
■ take 3 or more alcoholic drinks every day while using this product  
■ take other drugs containing aspirin or acetaminophen, or any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Drug Facts (continued)**

**Medication overdose headache warning:** Headaches may worsen if this product is used for 10 or more days per month. **Caffeine warning:** The recommended dose of the product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heartbeat.  
**Do not use**  
■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
**Ask a doctor before use if**  
■ you have never had a migraine diagnosed by a health professional  
■ you have a headache that is different from your usual migraines  
■ you have the worst headache of your life  
■ you have been and still are taking  
■ you have been taking aspirin or caused by head injury, operation, coughing or vomiting  
■ you have experienced your first headache after the age of 50  
■ you have daily headaches  
■ you have a migraine so severe as to require bed rest  
■ you have liver disease  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver disease, kidney disease, or asthma  
■ you have asthma  
■ you have problems or sensitive site effects from taking pain relievers or fever reducers  
■ you have vomiting with your migraine headache  
■ you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**

■ taking any prescription drug for:  
■ diabetes ■ gout ■ arthritis  
■ under a doctor's care for any serious condition  
■ taking any other drug  
■ taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Drug Facts (continued)**

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ your migraine is not relieved or worsens after first dose  
■ new or unexpected symptoms occur  
■ ringing in the ears or loss of hearing occurs  
**If pregnant or breast-feeding, ask a health or dental professional** before use. It's especially important to use aspirin at 20 weeks or later pregnancy unless a health professional has decided to do so by a doctor because it may cause problems in the unborn child or complication during delivery.  
**Keep out of reach of children.**  
In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222. Quick medical attention is critical for adults as well as for children and if you do not notice any signs or symptoms.

**Directions**

■ do not use more than directed  
■ adults: take 2 tablets with a glass of water  
■ if symptoms persist or worsen, ask your doctor  
■ do not take more than 2 tablets in 24 hours, unless directed by a doctor  
■ under 18 years of age: ask a doctor

**Other information**

■ store at 20°-25°C (68°-77°F) ■ dose cap tightly after use  
■ read all product information before using. Keep this book for important information.  
**Inactive ingredients:** black iron oxide, carnation wax, croscollonite, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, gelatin, glycerin, hypromellose, maltitol sorbitol, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polyvinyl alcohol, phthalate, polydioxane, pregelatinized starch, propylene glycol, stearic acid, titanium dioxide

**Questions or comments? 1-800-716-0990**

amazon **basic care**

Treats Migraine NDC 72288-295-27


**Migraine Relief**

Acetaminophen/Aspirin (NSAID)/Caffeine Tablets  
Pain Reliever/Pain Reliever Aid

Compare to Excedrin® Migraine active ingredients!

actual size 

80 GELTABS\*\* \*\*Gelatin Coated Tablets



Gluten Free  
Migraine  
This product is not  
manufactured or  
distributed by SK  
Consumer Health Care.  
distributed by SK  
© 2022 AMAZON.COM, INC. OR  
SEATTLE, WA 98109  
AMAZON.COM SERVICES LLC  
410 TERRY AVENUE N.  
DISTRIBUTED BY:  
AMAZON.COM SERVICES LLC  
DO NOT USE IF PRINTED FOIL UNDER CAP  
IS BROKEN OR MISSING  
KEEP BOX FOR REFERENCE



<b>BASIC CARE MIGRAINE RELIEF</b>			
acetaminophen, aspirin, caffeine tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72288-295
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FERROSOFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	GREEN, WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	M
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72288-295-27	1 in 1 CONTAINER	07/21/2022	
1		80 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075794	07/21/2022	

