## CHILDRENS MAPAP- acetaminophen tablet, chewable Major Pharmaceuticals

-----

Major 44-450-children

### Active ingredient (in each chewable tablet)

Acetaminophen 160 mg

### Purpose

Pain reliever/fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - sore throat
  - flu
  - toothache
  - the common cold
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### Ask a doctor before use if your child has

liver disease.

### Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

### Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

<sup>\*</sup>or as directed by a doctor

### Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- avoid high humidity
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### Inactive ingredients

citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

### Questions or comments?

1-800-426-9391

### **Principal Display Panel**

**MAJOR®** 

NDC 0904-6645-24 Compare to the active ingredient in Children's Tylenol® Chewables†

Children's **Māpap**®

# ACETAMINOPHEN 160 mg Chewable Tablets

Pain reliever / Fever reducer

ASPIRIN FREE • IBUPROFEN FREE

AGES 2-11

Chew or crush tablets completely before swallowing.

### 24 Chewable Tablets, 160 mg EACH

Actual Size

Bubble Gum Flavor

## TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING


Distributed by:

### MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268 (800) 616-2471 www.major-rugby.com Rev. 11/24 M-17 Re-order No. 700846

<sup>†</sup>This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Children's Tylenol® Chewables.

50844 REV0424E45008



Allergy alert: Acetaminophen may cause severe skin ■ with other drugs containing acetaminophen dainy amount Severe liver damage may occur if your child takes

more than 5 doses in 24 hours, which is the maximum

**Drug Facts** (continued)

Liver warning: This product contains acetaminophen. SprimbW

femborarijy reduces fever

■ toothache ■ the common cold ■ headache ■ sore throat ■ flu temporarily relieves minor aches and pains due to:

(in each chewable tablet) Active ingredient

IPLETE PRODUCT INFORMATIO

Drug Facts

səsŋ

**MAJOR®** 

NDC 0904-6645-24 Compare to the active ingredient in Children's Tylenol® Chewables®

# Children's *AMINOPHEN*

əsodınd

160 mg Chewable Tablets

Pain reliever / Fever reducer ASPIRIN FREE • IBUPROFEN FREE

AGES

Chew or crush tablets completely before swallowing

**24** Chewable Tablets, 160 mg EACH

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



Bubble Gum Flavor

B-1212-450-08-R REV0424E45008



BEV0424E45008 Children's Tylenol® Chewables. \*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark

Questions or comments?

Drug Facts (continued)

boj/etji/jeue' atestic acid, sucralose ethylcellulose, flavor, magnesium stearate, mannitol, citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated,

#### Inactive ingredients

- see end flap for expiration date and lot number

	4b41	
Stablets	ш	72-95
2 1/2 tablets	01-6	12-09
2 tablets	8-9	69-89
219lds/17 T	g-þ	74-85
feldst f	2-3	24-35
Ask a doctor	Under 2	Under 24
Dose (chewable tablets)*	(ry) agA	Weight (lb)

a do not give more than 5 times in № hours ■ repeat dose every 4 hours while symptoms last

Drug Facts (continued)

BLOOM STBLEET WOLLSWR TON OD cyew or crush tablets completely before swallowing;

away. Prompt medical attention is critical for adults as medical help or contact a Poison Control Center right

Ask a doctor or pharmacist before use if your child is

**Drug Facts** (continued)

weight to dose; otherwise, use age ■ find the right dose on chart below. If possible, use ■ do not give more than directed complete warnings for adult use this product does not contain directions or Directions well as for children even if you do not notice any signs or Keep out of reach of children. In case of overdose, get ■ sny new symptoms appear Tredness or swelling is present ■ tever gets worse or lasts more than 3 days ■ pain gets worse or lasts more than 5 days Stop use and ask a doctor if taking the blood thinning drug warfarin. Ask a doctor before use if your child has liver disease.

ask a doctor or pharmacist. are not sure whether a drug contains acetaminophen,

12°-30°C (59°-86°F) ■ store at 25°C (77°F); excursions permitted between PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ avoid high humidity REMPRE EVIDENT: DO NOT USE IF OUTRE noitemiotni ishto n ga ni acrin nà g nocini Do not use with any other drug containing scetaminophen (prescription or nonprescription). If you are not sure whether a drug contains season makes are not sure whether a drug contains and are not are not a contained to the contained to the

Distributed by:
MAJOR® PHARMACEUTICALS
Indianapolis, IN 46268
(800) 616-2471
www.major-rugby.com
Rev. 11/24 M-17 Re-order No. 700846

No Print Area Lot no. & Exp. Date

### **CHILDRENS MAPAP**

acetaminophen tablet, chewable

Product	Intorm	ation
FIUMULL		ation

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-6645
I I OGGET I YPE	110111111010	item code (source)	1120103010013

**Route of Administration** ORAL

### **Active Ingredient/Active Moiety**

Ingredient Name	<b>Basis of Strength</b>	Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) ACETAMINOPHEN 160 mg

### **Inactive Ingredients**

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
D&C RED NO. 30 ALUMINUM LAKE (UNII: GE75M6YV5W)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 30WL53L36A)	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

STEARIC ACID (UNII: 4ELV7Z65AP)

SUCRALOSE (UNII: 96K6UQ3ZD4)

### **Product Characteristics**

Color	pink	Score	2 pieces
Shape	ROUND	Size	16mm
Flavor	BUBBLE GUM	Imprint Code	44;450
Contains			

Contains

### **Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6645- 24	4 in 1 CARTON	02/25/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### **Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/25/2005	

### Labeler - Major Pharmaceuticals (191427277)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0904-6645)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0904-6645) , pack(0904-6645)

Establishment						
Name	Address	ID/FEI	<b>Business Operations</b>			
LNK International, Inc.		967626305	pack(0904-6645)			

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(0904-6645)

Revised: 10/2025 Major Pharmaceuticals