

**ALAHIST D- pheniramine maleate, phenylephrine hcl tablet**  
**Poly Pharmaceuticals, Inc.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**Alahist D**

**ACTIVE INGREDIENT**

**Active Ingredient (in each tablet) Purpose**

---

Pheniramine Maleate 17mg	Antihistamine
Phenylephrine HCl 10mg	Nasal Decongestant

---

**USES**

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- ☐ runny nose
- ☐ sneezing
- ☐ itching of the nose or throat
- ☐ itchy, watery eyes
- ☐ nasal congestion ☐
  - reduces swelling of nasal passages

**WARNINGS**

**Do not exceed recommended dosage.**

**Do not use this product**

☐ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- ☐ a breathing problem such as emphysema or chronic bronchitis
- ☐ glaucoma
- ☐ trouble urinating due to enlargement of the prostate gland
- ☐ heart disease
- ☐ high blood pressure
- ☐ thyroid disease ☐
- diabetes

**Ask a doctor or pharmacist before use if you** are taking sedatives or tranquilizers.

### **When using this product**

- excitability may occur, especially in children
- may cause drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever

**If pregnant or breast-feeding, ask a health professional before use.**

## **DIRECTIONS**

### **Directions**

---

Adults and children 12 years of age and over	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours
Children 6 to under 12 years of age	1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours
Children under 6 consult a doctor	

---

## **INACTIVE INGREDIENTS**

### **Inactive ingredients**

Magnesium Stearate, Microcrystalline Cellulose, Natural Yellow, Sodium Starch Glycolate

## **QUESTIONS**

Questions? Comments?

Call 1-800-882-1041

Manufactured for:

Poly Pharmaceuticals

Huntsville, AL 35763 Rev. 04/17

## **OTHER INFORMATION**

### **Other information**

Store at 15°-30°C (59°-86°F). Supplied in a tight, light-resistant container with a child-resistant cap. Alahist D Tablets are yellow, debossed "A" bisect "D" on one side and plain on the other.

### **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

Antihistamine

# Nasal Decongestant

## Package Label



<p><b>Lot:</b> 50991-776-01-0</p> <p><b>NDC 50991-776-01</b></p> <p><b>ALAHIST® D Tablets</b></p> <p>Antihistamine • Nasal Decongestant</p> <p>Each tablet contains: Pheniramine Maleate ..... 17.5 mg Phenylephrine Hydrochloride ... 10 mg</p> <p>100 Tablets</p>	<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each tablet)</b></p> <p>Pheniramine Maleate 17.5 mg ..... Antihistamine Phenylephrine Hydrochloride 10 mg ..... Nasal Decongestant</p> <p><b>Purpose</b></p> <p><b>Uses</b></p> <p>Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:</p> <ul style="list-style-type: none"> <li>runny nose</li> <li>sneezing</li> <li>itching of the nose or throat</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>itchy, watery eyes</li> <li>nasal congestion</li> <li>reduces swelling of nasal passages</li> </ul> <p><b>Warnings</b></p> <p>Do not exceed recommended dosage.</p> <p>Do not use this product</p> <ul style="list-style-type: none"> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p>disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>a breathing problem such as emphysema or chronic bronchitis</li> <li>glaucoma</li> <li>trouble urinating due to enlargement of the prostate gland</li> <li>heart disease</li> <li>high blood pressure</li> <li>thyroid disease</li> <li>diabetes</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.</b></p> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>excitability may occur, especially in children</li> <li>may cause drowsiness</li> <li>avoid alcoholic drinks</li> <li>alcohol, sedatives, and tranquilizers may increase the drowsiness effect</li> <li>use caution when driving a motor vehicle or operating machinery</li> </ul>						
11.8438"										
<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>nervousness, dizziness, or sleeplessness occur</li> <li>symptoms do not improve within 7 days or are accompanied by fever</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <table border="1"> <tr> <td>Adults and children 12 years of age and over:</td> <td>1 tablet every 4 hours not to exceed 6 tablets in 24 hours or as directed by a doctor</td> </tr> <tr> <td>Children 6 to under 12 years of age:</td> <td>1/2 tablet every 4 hours not to exceed 3 tablets in 24 hours or as directed by a doctor</td> </tr> <tr> <td>Children under 6 years of age:</td> <td>Consult a doctor</td> </tr> </table>	Adults and children 12 years of age and over:	1 tablet every 4 hours not to exceed 6 tablets in 24 hours or as directed by a doctor	Children 6 to under 12 years of age:	1/2 tablet every 4 hours not to exceed 3 tablets in 24 hours or as directed by a doctor	Children under 6 years of age:	Consult a doctor	<p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <p>Store at 15°-30°C (59°-86°F). Supplied in a tight, light-resistant container with a child-resistant cap. Alahist D Tablets are yellow tablets debossed A   D on one side and plain on the other side.</p> <p><b>Inactive ingredients</b></p> <p>Magnesium stearate, microcrystalline cellulose, natural yellow, sodium starch glycolate</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Questions? Comments?</b></p> <p>Call 1-800-882-1041</p> <p>Manufactured for: Poly Pharmaceuticals Huntsville, AL 35763</p> <p>Rev. 01/20</p>	
Adults and children 12 years of age and over:	1 tablet every 4 hours not to exceed 6 tablets in 24 hours or as directed by a doctor									
Children 6 to under 12 years of age:	1/2 tablet every 4 hours not to exceed 3 tablets in 24 hours or as directed by a doctor									
Children under 6 years of age:	Consult a doctor									

## ALAHIST D

pheniramine maleate, phenylephrine hcl tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50991-776
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENIRAMINE MALEATE</b> (UNII: NYW905655B) (PHENIRAMINE - UNII:134FM9ZZ6M)	PHENIRAMINE MALEATE	17.5 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

## Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	

## Product Characteristics

Color	yellow	Score	2 pieces
Shape	OVAL	Size	11mm
Flavor		Imprint Code	A;D
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50991-776-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2020	
2	NDC:50991-776-02	12 in 1 BLISTER PACK; Type 0: Not a Combination Product	04/01/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/01/2020	

**Labeler** - Poly Pharmaceuticals, Inc. (198449894)

Revised: 1/2024

Poly Pharmaceuticals, Inc.