

ASPIRIN- aspirin tablet
HART Health

ASPIRIN

Active Ingredient

Aspirin (NSAID)* 325mg (*non-steroidal anti-inflammatory drug)

Purpose

Pain Reliever/Fever Reducer

Uses

Temporarily relieves minor aches and pains due to:

- headache
- backache
- muscular aches
- minor arthritis pain
- common cold
- toothache
- menstrual cramps

Temporarily reduces fever

Warnings

Reye's Syndrome: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin

Allergy alert: Aspirin may cause a several allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, others)
- have 3 or more alcoholic drinks every day while using this product

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have asthma, high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you:

- take any other drug containing an NSAID
- take a blood thinning (anticoagulant) or steroid drug
- take any drug for diabetes, gout, or arthritis

- an allergic reaction occurs, seek medical help right away
- fever gets worse or lasts more than 3 days
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- ringing in the ears or loss of hearing occurs
- new symptoms occur
- you have any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, or stomach pain that does not get better

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions:

Do not take more than directed.


Adults and children 12 years of age and over:

- take 1 or 2 tablets with water every 4 hours while symptoms persist
- do not take more than 12 tablets in 24 hours unless directed by a doctor

Children under 12 years of age: ask a doctor

Inactive Ingredients: corn starch, hypromellose, polyethylene glycol, propylene glycol

50 count
 PMS 279C blue
 PMS 7404C yellow
 small medication box dieline
 10.8cm x 6cm x 9cm



NDC 50332-0139-4

Item 5521

Item 5521

COMPARE TO
Bayer®
 Active Ingredient

COMPARE TO
Bayer®
 Active Ingredient

Aspirin
Pain Reliever

Aspirin
Pain Reliever
Analgésico

Aspirin (NSAID)
 Reduces Pain and Fever

Aspirin (NSAID)
 Reduces Pain and Fever
 Reduce el Dolor y la Fiebre

50 packets of 2 tablets
 100 total tablets

50 packets of 2 tablets
 100 total tablets

This package for households
 without young children.

This package for households
 without young children.

Manufacturer for:
HARTHealth
 Seattle, WA 98124 www.harthealth.com



*This product is not manufactured or distributed by
 Bayer HealthCare LLC, owner of the registered
 trademark Bayer® products.

Drug Facts

Aspirin (NSAID) 325 mg (nonsteroidal anti-inflammatory drug) Pain reliever/fever reducer

Uses: Temporarily relieve minor aches and pains due to: ■ headache ■ muscle aches ■ backache ■ minor arthritis pain ■ common cold ■ toothache ■ menstrual cramps Temporarily reduce fever

Warnings: **Reye's syndrome:** Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

Warnings: Aspirin may cause a severe allergic reaction which may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ have 3 or more alcoholic drinks every day while using this product ■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen or others)

Do not use: If you have ever had an allergic reaction to any pain reliever/fever reducer.

Ask a doctor before use if:
 ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn ■ you have asthma, high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have a history of stomach bleeding, such as heartburn ■ you have a history of stomach problems, such as heartburn ■ you have a history of stomach bleeding, such as heartburn ■ you have a history of stomach bleeding, such as heartburn

Ask a doctor or pharmacist before use if you:
 ■ take any drug containing an NSAID ■ take any drug for diabetes, gout, or arthritis ■ take any drug for diabetes, gout, or arthritis ■ take any drug for diabetes, gout, or arthritis

Stop use and ask a doctor if:
 ■ fever gets worse or lasts more than 3 days ■ stomach bleeding warning applies to you ■ pain gets worse or lasts more than 10 days ■ new symptoms occur ■ an allergic reaction occurs, seek medical help right away ■ stomach pain that does not get better ■ ringing of the ears or loss of hearing occurs

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222


Directions
 Do not take more than directed.
 Adults and children 12 years of age and over:
 ■ take 1 or 2 tablets with water every 4 hours while symptoms persist.
 ■ Do not take more than 12 tablets in 24 hours unless directed by a doctor.
 Children under 12 years of age: ask a doctor.

Other information
 ■ tablets enclosed in a sealed packet
 ■ do not use if packet is torn, out, or open
 ■ store at 59° to 86°F (15°-30°C)
 ■ avoid excessive heat and humidity

Inactive ingredients
 corn starch, hypromellose, polyethylene glycol, propylene glycol

KEY

- cut line —————
- inward fold line - - - - -
- outward fold line ······
- perf ————
- slit ————
- outside glue
- inside glue



50 count
 PMS 279C blue
 PMS 7404C yellow
 small medication box dieline
 10.8cm x 6cm x 9cm

NDC 50332-0139-4

Item 5521

Item 5521

Aspirin
Pain Reliever

Aspirin
Pain Reliever
Analgésico

Aspirin (NSAID)
 Reduces Pain and Fever

Aspirin (NSAID)
 Reduces Pain and Fever
 Reduce el Dolor y la Fiebre

Drug Facts (continued)

Aspirin
Pain Reliever
Aspirin (NSAID)

Active Ingredient (in each tablet)
 Aspirin (acetylsalicylic acid) 325 mg (non-steroidal anti-inflammatory drug)

Purpose
 Pain reliever/fever reducer

Warnings **Ryze's syndrome:** Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Ryze's syndrome, a rare but serious illness reported to be associated with aspirin.

Other warnings: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Ryze's syndrome, a rare but serious illness reported to be associated with aspirin.

Other warnings: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Ryze's syndrome, a rare but serious illness reported to be associated with aspirin.

Other warnings: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Ryze's syndrome, a rare but serious illness reported to be associated with aspirin.

Drug Facts (continued)

Directions

Do not take more than directed.

Adults and children 12 years of age and over:

- Take 1 or 2 tablets with water every 4 hours while symptoms persist.
- Do not take more than 12 tablets in 24 hours unless directed by a doctor.

Children under 12 years of age: ask a doctor.

Other information

- Tablets enclosed in a sealed packet.
- Do not use if packet is torn, out, or open.
- Store at 59° to 86°F (15°-30°C).
- Avoid excessive heat and humidity.

Inactive ingredients

corn starch, hydroxypropylcellulose, polyethylene glycol, propylene glycol

Manufacturer for:
HART-Health
 Seattle, WA 98124 www.hart-health.com
 9471-23

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® products.

KEY

- cut line _____
- inward fold line - - - - -
- outward fold line ·····
- perf - - - - -
- slit - - - - -
- outside glue
- inside glue

ASPIRIN
 aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50332-0139
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWJLA)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASPIRIN;44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50332-0139-4	50 in 1 BOX, UNIT-DOSE	01/01/2023	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:50332-0139-7	125 in 1 BOX, UNIT-DOSE	01/01/2023	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:50332-0139-8	250 in 1 BOX, UNIT-DOSE	01/01/2023	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:50332-0139-3	25 in 1 BOX, UNIT-DOSE	01/01/2023	
4		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M	01/01/2023	

Labeler - HART Health (069560969)

Revised: 1/2026

HART Health