

LEADER ANTI-DIARRHEAL- loperamide hcl tablet, film coated
Cardinal Health

Cardinal Health Anti-Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are

taking antibiotics

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25° C (68°-77° F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-800-719-9260

Principal Display Panel

Bonus Size

Compare to Imodium® A-D active ingredient

SEE NEW WARNINGS AND DIRECTIONS

Loperamide Hydrochloride Tablets, 2mg

Anti-Diarrheal

Anti-Diarrheal

Controls the Symptoms of Diarrhea

Actual Size

SATISFACTION GUARANTEED

18 CAPLETS*

12 + 6 FREE = 18 CAPLETS

*(capsule-shaped tablets)

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

NDC 49761-128-18

Bonus Size

LEADER®

Compare to **Imodium® A-D**
active ingredients

SEE NEW WARNING AND DIRECTIONS

Loperamide Hydrochloride Tablets, 2mg

Anti-Diarrheal

Anti-Diarrheal

Controls the Symptoms of Diarrhea

A2054E9C2



18 CAPLETS*

12 + 6 FREE = 18 CAPLETS

Actual Size



(*capsule-shaped tablets)

A2054 E9 C2

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium

Drug Facts

Active Ingredient (in each caplet) Loperamide HCl 2 mg **Purpose** Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
 - diarrhea lasts for more than 2 days
 - you get a abdominal swelling or bulging.
- These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of

Drug Facts (continued)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use a ge.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other Information


- store at 20°-25° C (68°-77° F)

All Leader® Brand products are 100% satisfaction guaranteed or return to place of purchase for a full refund.

DISTRIBUTED BY
CARDINAL HEALTH
DUBLIN, OHIO 43017
CIN 501 3347
www.myleader.com
1-800-200-6313



6295 12638 9



LEADER ANTI-DIARRHEAL

loperamide hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49781-128
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:49781-128-18	18 in 1 CARTON	09/25/2014	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA075232		09/25/2014	

Labeler - Cardinal Health (097537435)

Revised: 11/2019

Cardinal Health