

**IBUPROFEN IMMEDIATE RELEASE- ibuprofen tablet, coated**  
**Discount Drug Mart**

-----  
**Ibuprofen tablets USP 200mg**  
**Discount Drug Mart**

**Drug Facts**

**ACTIVE INGREDIENT(S)**

*(in each brown tablet)*

Ibuprofen USP 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**PURPOSE**

Pain reliever/fever reducer

**USE(S)**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**WARNINGS**

**Allergy alert**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **DO NOT USE**

- if you have ever had an allergic reaction to any other pain reliever / fever reducer
- right before or after heart surgery

### **ASK A DOCTOR BEFORE USE IF**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

### **ASK A DOCTOR OR PHARMACIST BEFORE USE IF**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### **WHEN USING THIS PRODUCT**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### **STOP USE AND ASK DOCTOR IF**

- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a poison control center (1-800-222-1222) right away.

## DIRECTIONS

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	<ul style="list-style-type: none"><li>• take 1 tablet every 4 to 6 hours while symptoms persist</li><li>• if pain or fever does not respond to 1 tablet, 2 tablets may be used</li><li>• do not exceed 6 tablets in 24 hours, unless directed by a doctor</li></ul>
children under 12 years	<ul style="list-style-type: none"><li>• ask a doctor</li></ul>

## OTHER INFORMATION

- store between 20 – 25°C (68-77°F)
- **tamper evident:** do not use if imprinted safety seal under cap is broken or missing
- see end panel for lot number and expiry date

## INACTIVE INGREDIENT (S)

colloidal silicon dioxide, corn starch, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, red iron oxide, sodium starch glycolate, titanium dioxide, triacetin and talc

## QUESTIONS OR COMMENTS?

Adverse drug event Call: (877) 586-7979

## PRINCIPAL DISPLAY PANEL

Package Label (Round Shaped Tablets) - Principal Display Panel - 50s - Count Carton, 200 mg Tablets

DISCOUNT

drug mart

FOOD FAIR

NDC 53943-401-09

*SEE NEW WARNINGS INFORMATION*

*IBUPROFEN*

TABLETS, USP 200 mg

Pain Reliever / Fever Reducer (NSAID)\*

IBU

200

*50 COATED TABLETS*

\*nonsteroidal anti-inflammatory drug

Compare to the Active Ingredient of ADVIL<sup>®</sup>\*\*

\*\*This product is not manufactured or distributed by Pfizer Inc. owner of the registered trade mark Advil<sup>®</sup>

<b>IBUPROFEN IMMEDIATE RELEASE</b>			
ibuprofen tablet, coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53943-401

Open From This Side

LOT: \_\_\_\_\_  
EXP: \_\_\_\_\_

020002795

9 335511017777 0

Important: Read all product information before using.  
Keep this box for important information.

**Drug Facts (Continued)**

- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**  
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- ask a doctor

children under 12 years

**Other Information**

- Store between 20-25°C (68-77°F)
- tamper evident**: do not use if imprinted safety seal under cap is broken or missing
- see end panel for lot number and expiry date

DISCOUNT **drug mart** FOOD & DRUG STORE

NDC 53943-401-09

**SEE NEW WARNINGS INFORMATION**

**IBUPROFEN**

TABLETS, USP 200 mg

Pain Reliever / Fever Reducer (NSAID)\*

50 COATED TABLETS

IBU 200

\*nonsteroidal anti-inflammatory drug

Compare to the Active Ingredient of ADVIL\*\*

\*\*This product is not manufactured or distributed by Pfizer Inc. owner of the registered trade mark Advil®

020002795

**Drug Facts**

**Active ingredient (in each brown tablet)**  
Ibuprofen USP 200 mg (NSAID)\* ..... Pain reliever/fever reducer

\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - muscular aches
  - toothache
  - the common cold
- headache
- minor pain of arthritis
- backache
- menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- rash
- blisters
- shock
- skin redness
- asthma (wheezing)

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (Continued)**

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever / fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days

43x43x78mm

**Drug Facts (Continued)**

**Inactive Ingredients**

colloidal silicon dioxide, corn starch, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, red iron oxide, sodium starch glycolate, titanium dioxide, triacetin and talc

**Questions or comments?**  
Advise drug event Call: (877) 586-7979

SAFETY INFORMATION: IF YOU PURCHASE THIS PRODUCT FROM A PHARMACY, DISCOUNT DRUG MART, FOOD & DRUG STORE, OR OTHER RETAILER, PLEASE RETURN THE PRODUCT TO THE STORE OF PURCHASE. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISCOUNT DRUG MART 211 COMMERCIAL DRIVE, MEDINA, OHIO 44256

Mg Lic No.: DRUGSP/05 13 1523  
Distributed by: Drug Mart - Food & Drug Mart  
Medina, Ohio 44256. www.discount-drugmart.com

<b>Route of Administration</b>	ORAL
--------------------------------	------

<b>Active Ingredient/Active Moiety</b>		
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

<b>Inactive Ingredients</b>	
<b>Ingredient Name</b>	<b>Strength</b>
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
HYPROMELLOSE 2910 (6 MPAS) (UNII: 0WZ8WG20P6)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TALC (UNII: 7SEV7J4R1U)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

<b>Product Characteristics</b>			
<b>Color</b>	BROWN	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND SHAPED)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	IBU200
<b>Contains</b>			

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-401-09	1 in 1 CARTON		
1		50 in 1 BOTTLE		
2	NDC:53943-401-10	1 in 1 CARTON		
2		100 in 1 BOTTLE		
3	NDC:53943-401-15	500 in 1 BOTTLE		

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079129	12/22/2014	

**Labeler** - Discount Drug Mart (047741335)

**Registrant** - Shasun Pharmaceuticals Limited (915786829)

## Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	MANUFACTURE(53943-401)

Revised: 11/2014

Discount Drug Mart